arXiv:math/0410590v1 [math.GR] 27 Oct 2004

PRODUCTS OF CHARACTERS WITH FEW IRREDUCIBLE CONSTITUENTS

EDITH ADAN-BANTE

Abstract. We study the solvable groups G that have an irreducible character $\chi \in \mathrm{Irr}(G)$ such that $\chi\overline{\chi}$ has at most two non-principal irreducible constituents.

1. Introduction

Let G be a finite group. Denote by $\mathrm{Irr}(G)$ the set of irreducible complex characters of G. Let 1_G be the principal character of G. Denote by $[\Theta, \Phi]$ the inner product of the characters Θ and Φ of G. Through this work, we will use the notation of [5].

Let $\chi \in \mathrm{Irr}(G)$. Define $\overline{\chi}(g)$ to be the complex conjugate $\overline{\chi(g)}$ of $\chi(g)$ for all $g \in G$. Then $\overline{\chi}$ is also an irreducible complex character of G. Since the product of characters is a character, then $\chi\overline{\chi}$ is a character of G. So it can be expressed as an integral linear combination of irreducible characters. Now observe that

$$[\chi\overline{\chi}, 1_G] = [\chi, \chi] = 1$$

where the last equality holds since $\chi \in \mathrm{Irr}(G)$. Assume now that $\chi(1) > 1$. Then the decomposition of the character $\chi\overline{\chi}$ into its irreducible constituents $\alpha_1,\ \alpha_2, \ldots, \alpha_n$ has the form

$$\chi\overline{\chi} = 1_G + \sum_{i=1}^{n} m_i \alpha_i$$

where $m_i > 0$ is the multiplicity of α_i. Set $\eta(\chi) = n$. Thus $\eta(\chi)$ is the number of distinct non-principal irreducible constituents of $\chi\overline{\chi}$.

In Theorem A of [1], it is proved that there exist universal constants C and D such that for any finite solvable group G and any $\chi \in \mathrm{Irr}(G)$, we have that $\mathrm{dl}(G/\mathrm{Ker}(\chi)) \leq C\eta(\chi\overline{\chi}) + D$. In Theorem B of [1], it is proved that if G is a finite solvable group and $\chi \in \mathrm{Irr}(G)$, then $\chi(1)$ has at most $\eta(\chi)$ distinct prime divisors. If, in addition, G is supersolvable, then $\chi(1)$ is the product of at most $\eta(\chi) - 1$ primes. The purpose of this note is to study the solvable groups G that have faithful character $\chi \in \mathrm{Irr}(G)$ such that $\eta(\chi) \leq 2$.

Let G be a solvable group and $\chi \in \mathrm{Irr}(G)$ be a faithful character. Assume that $\eta(\chi) = 1$, i.e $\chi\overline{\chi} = 1_G + m\alpha$ for some $\alpha \in \mathrm{Irr}(G)$ and some integer $m > 0$. Since $\chi\overline{\chi}$ and 1_G are real characters, so is α. Thus G has an irreducible real character. Therefore G has even order. More can be said in that regard and Theorem A "classifies" those groups.

Date: 2004.

1991 *Mathematics Subject Classification.* 20c15.

Key words and phrases. Product of characters, irreducible character, derived length.

This research was partially supported by grant NSF DMS 9970030.

Theorem A. *Let G be a finite solvable group and $\chi \in \mathrm{Irr}(G)$ be a faithful character. Assume that*

$$\chi\overline{\chi} = 1_G + m\alpha \tag{1.0.1}$$

for some $\alpha \in \mathrm{Irr}(G)$ and some positive integer m. Set $Z = \mathbf{Z}(G)$. Let E/Z be a chief factor of G. Set $\bar{G} = G/E$. Then $m = 1$, $\chi(1)^2 = |E : Z|$ and one of the following holds:

(i) $\chi(1) = 2$ and $G/Z \cong A_4$ or $G/Z \cong S_4$.

(ii) $\chi(1) = 3$ and either $\bar{G} \cong \mathbf{Q}_8$ or $\bar{G} \cong \mathrm{SL}(2,3)$.

(iii) $\chi(1) = 5$ and $\bar{G} \cong \mathrm{SL}(2,3)$.

(iv) $\chi(1) = 7$ and $\bar{G} \cong \widetilde{\mathrm{GL}}(2,3)$.

(v) $\chi(1) = 9$, $\mathbf{F}(\bar{G})$ is an extra-special group of order 2^5, namely the central product of the quaternion group $\mathbf{Q}_8$ and the dihedral group of order 8, $|\mathbf{F}_2(\bar{G})/\mathbf{F}(\bar{G})| = 5$ and $\bar{G}/\mathbf{F}_2(\bar{G})$ is a subgroup of the cyclic group of order 2.

We can check that in all the cases given by Theorem A we have that $|G : Z| \leq 25920$. Assume now that G is a solvable group, $\chi \in \mathrm{Irr}(G)$ is a faithful character and $\eta(\chi) = 2$. In such situation, the index of the center Z of G in G is not longer bounded by a universal constant, not even if in addition the group is nilpotent (see Subsection 4.4). But the derived length $\mathrm{dl}(G)$ is at most 18 and we have more information regarding the structure of the group G.

Theorem B. *Let G be a finite solvable group and $\chi \in \mathrm{Irr}(G)$ be a faithful character. Assume*

$$\chi\overline{\chi} = 1_G + m_1\alpha_1 + m_2\alpha_2$$

where $\alpha_1, \alpha_2 \in \mathrm{Irr}(G)$ are non-principal characters, m_1 and m_2 are strictly positive integers. Let Z be the center of G. Then

(i) The order of G is even.

(ii) $\mathrm{dl}(G) \leq 18$.

(iii) Either $\mathrm{Ker}(\alpha_1) = Z$ or $\mathrm{Ker}(\alpha_2) = Z$. If $\mathrm{Ker}(\alpha_1)$ and $\mathrm{Ker}(\alpha_2)$ are both abelian subgroups, then $\chi(1)$ is a power of a prime number. Otherwise $\{m_1, m_2\} = \{1, 4\}$ and $\chi(1) \in \{10, 14\}$.

Propositions 4.2.5, 4.2.9 and 4.2.13 describe with more detail the possible structure of the group G. Examples of groups and characters satisfying the hypotheses of Theorem B are studied in Subsection 4.3.

Acknowledgment. This is part of my Ph.D. Thesis. I thank Professor Everett C. Dade, my adviser, for his advise, suggestions and generosity with his time. I thank Professor I. Martin Isaacs for suggesting the question that started my studies in products of characters. I would also like to thank the mathematics department of the University of Wisconsin, at Madison, for their hospitality while I was visiting Professor Isaacs, and the mathematics department of the University of Illinois at Urbana-Champaign for their support.

2. VECTOR SPACES

B. Huppert classified all the solvable groups that act faithfully on a finite vector space V and transitively on $V^{\#} = V \setminus \{0\}$ (see Lemma 2.1.2). If in addition V is a symplectic vector space and the action of G on V preserves the symplectic form, the structure of the group G must be very constrained. We will classify the solvable groups G such that G can act faithfully on a symplectic vector space preserving the symplectic form and acting transitively on $V^{\#}$.

In Subsection 2.1 we mention the results that we use in the proof of Theorem 2.2.1, the main result of this section.

2.1. **Preliminaries.** Let V be a vector space of dimension n over $\mathrm{GF}(q)$. We may assume that V is $\mathrm{GF}(q^n)$, the Galois field with q^n elements, since $\mathrm{GF}(q^n)$ is a vector space of dimension n over $\mathrm{GF}(q)$. Let $\mathcal{G} = \mathrm{Gal}(\mathrm{GF}(q^n)/\,\mathrm{GF}(q))$ be the group of automorphism of the field $\mathrm{GF}(q^n)$ fixing $\mathrm{GF}(q)$. Then the *semi-linear group* $\Gamma(V)$ of V is

$$\Gamma(V) = \{x \mapsto ax^{\sigma} \ |a \in (\mathrm{GF}(q^n))^{\#}, \sigma \in \mathcal{G}\}$$

Observe that

$$U = \{x \mapsto ax \ |a \in (\mathrm{GF}(q^n))^{\#}\} \tag{2.1.1}$$

is a normal subgroup of $\Gamma(V)$. Also observe that $U \cong (\mathrm{GF}(q^n))^{\#}$, the group of units of the field $\mathrm{GF}(q^n)$. So U is a cyclic subgroup of $\Gamma(V)$ of order $q^n - 1$. Recall that $\mathcal{G}$ is a cyclic group of order n. Observe that $\Gamma(V)/U \cong \mathcal{G}$. Therefore $\Gamma(V)/U$ is a cyclic group of order n.

Lemma 2.1.2 (B. Huppert)**.** *Let V be a vector space of dimension n over* $\mathrm{GF}(q)$, *where q is a prime power. Suppose that G is a solvable subgroup of $GL(V)$ that transitively permutes the elements of $V^{\#}$. Let $\mathbf{F}(G)$ be the Fitting subgroup of G. Then one of the following holds:*

(i) G is isomorphic to a subgroup of $\Gamma(V)$.

(ii) $q^n = 3^4$, $\mathbf{F}(G)$ is extra-special of order 2^5, $|\mathbf{F}_2(G)/\mathbf{F}(G)| = 5$ and $G/\mathbf{F}_2(G) \leq \mathbf{Z}_4$.

(iii) $q^n = 3^2$, 5^2, 7^2, 11^2 or 23^2. Here $\mathbf{F}(G) = QT$, where $T = \mathbf{Z}(G) \leq \mathbf{Z}(GL(V))$ is cyclic, $\mathbf{Q}_8 \cong Q \trianglelefteq G$, $T \cap Q = \mathbf{Z}(Q)$ and $Q/Z(Q) \cong \mathbf{F}(G)/T$ is a faithful irreducible $G/\mathbf{F}(G)$-module. We also have one of the following entries:

q^n	$\lvert T\rvert$	$G/\mathbf{F}(G)$
3^2	*2*	$\mathbf{Z}_3$ *or* S_3
5^2	*2 or 4*	$\mathbf{Z}_3$
5^2	4	S_3
7^2	*2 or 6*	S_3
11^2	10	$\mathbf{Z}_3$ *or* S_3
23^2	22	S_3

Proof. See Theorem 6.8 of [9]. □

Let V be a finite dimensional vector space over the field F. We say that V is a *symplectic vector space* if there exists a non-degenerate, alternating and bilinear form f on V.

We denote by $\mathrm{Sp}(V)$ the group of automorphism of V preserving the symplectic form f. We denote by $\mathrm{Sp}(n, q)$ the group of automorphism of a n-dimensional symplectic vector space over $\mathrm{GF}(q)$. The following is a list of well known results that we will use later.

Lemma 2.1.3. *Let* $\mathrm{Sp}(2,q)$ *be the symplectic group where* $(q,2)=1$. *Then*

(i) $\mathrm{Sp}(2,q)=\mathrm{SL}(2,q)$.

(ii) $|\mathbf{Z}(\mathrm{SL}(2,q))|=2$.

(iii) If $q>3$, *then* $\mathrm{SL}(2,q)$ *is not a solvable group.*

(iv) $|\,\mathrm{SL}(2,q)|=q(q^2-1)$.

(v) $\mathrm{SL}(2,3)\cong \mathbf{Q}_8 \rtimes \mathbf{Z}_3$.

(vi) $\mathrm{SL}(2,3)$ *is isomorphic to a subgroup of* $\mathrm{SL}(2,5)$.

(vii) The 2-Sylow subgroups of $\mathrm{GL}(2,3)$ *are isomorphic to the semi-dihedral group of order* 16.

(viii) For q *odd, the 2-Sylow subgroups of* $\mathrm{SL}(2,q)$ *are isomorphic to generalized quaternion groups.*

Definition 2.1.4. We define $\widetilde{\mathrm{GL}}(2,3)$ as the group given by the following representation

$$< x,y,z \mid x^8=1,\ y^2=x^4,\ z^3=1,\ x^y=x^{-1},\ z^x=x^2yz^{-1},\ (x^2)^z=y > .$$

Lemma 2.1.5. $\widetilde{\mathrm{GL}}(2,3)$ *is not isomorphic to* $\mathrm{GL}(2,3)$. *Also* $\widetilde{\mathrm{GL}}(2,3)$ *has a subgroup of index 2 isomorphic to* $\mathrm{SL}(2,3)$, *and* $\mathbf{F}(\widetilde{\mathrm{GL}}(2,3))\cong\mathbf{Q}_8$.

Proof. We can check that the subgroup generated by x and y is a 2-Sylow subgroup of $\widetilde{\mathrm{GL}}(2,3)$ and is isomorphic to $\mathbf{Q}_{16}$. Thus $\mathrm{GL}(2,3)$ and $\widetilde{\mathrm{GL}}(2,3)$ are not isomorphic since a 2-Sylow subgroup of $\mathrm{GL}(2,3)$ is isomorphic to the semi-dihedral group of order 16.

Observe that $\mathbf{F}(\widetilde{\mathrm{GL}}(2,3))=\mathbf{Q}_8$, namely the subgroup generated by x^2 and y. Also observe that the subgroup generated by x^2, y and z is isomorphic to $\mathrm{SL}(2,3)$. □

The group $\widetilde{\mathrm{GL}}(2,3)$ is the isoclinic group of $\mathrm{GL}(2,3)$.

Lemma 2.1.6. *The group* $\mathrm{SL}(2,7)$ *has a subgroup isomorphic to* $\widetilde{\mathrm{GL}}(2,3)$. *Also, any subgroup of order 48 in* $\mathrm{SL}(2,7)$ *is isomorphic to* $\widetilde{\mathrm{GL}}(2,3)$.

Proof. Let $x,y,z\in\mathrm{SL}(2,7)$ be the matrices

$$x=\begin{pmatrix}2&2\\5&2\end{pmatrix},\quad y=\begin{pmatrix}3&2\\2&4\end{pmatrix},\quad z=\begin{pmatrix}4&0\\1&2\end{pmatrix}.$$

We can check that they satisfy the relations given in Definition 2.1.4. Thus $\mathrm{SL}(2,7)$ has a subgroup isomorphic to $\widetilde{\mathrm{GL}}(2,3)$, mainly the group generated by x, y and z.

Let H be a subgroup of $\mathrm{SL}(2,7)$ of order 48. Then H is a maximal subgroup of $\mathrm{SL}(2,7)$. We can check using the Atlas that the maximal subgroups of $\mathrm{SL}(2,7)$ of order 48 are all isomorphic. □

2.2. **Transitive actions and symplectic vector spaces.**

Theorem 2.2.1. *Let* G *be a finite solvable group. Let* V *be a symplectic vector space of dimension* n *over* $\mathrm{GF}(q)$, *where* q *is a power of a prime number. Assume that* V *is a faithful* G*-module and the action of* G *on* V *preserves the symplectic form. Also assume that* G *acts transitively on* $V^{\#}$. *Set* $e^2=|V|=q^n$. *Let* $\mathbf{F}(G)$ *be the Fitting subgroup of* G *and* $\mathbf{F}_2(G)/\mathbf{F}(G)$ *be the Fitting subgroup of* $G/\mathbf{F}(G)$. *Then one of the following holds:*

(i) $e=2$ *and* $G\cong\mathbf{Z}_3$ *or* $G\cong S_3$.

(ii) $e=3$, *and either* $G\cong\mathbf{Q}_8$ *or* $G\cong\mathrm{SL}(2,3)$.

(iii) $e = 5$ *and* $G \cong \mathrm{SL}(2,3)$.

(iv) $e = 7$ *and* $G \cong \widetilde{\mathrm{GL}}(2,3)$.

(v) $e = 9$, $\mathbf{F}(G)$ *is an extra-special group of order* 2^5, *namely the central product of the quaternion group* $\mathbf{Q}_8$ *and the dihedral group of order 8,* $|\mathbf{F}_2(G)/\mathbf{F}(G)| = 5$ *and* $G/\mathbf{F}_2(G)$ *is a subgroup of the cyclic group of order 2.*

Observe that the hypothesis of Theorem 2.2.1 implies the hypothesis of Lemma 2.1.2. Thus either (i), (ii) or (iii) of Lemma 2.1.2 holds. We will analyze each case given by the conclusion of Lemma 2.1.2.

Lemma 2.2.2. *Let* C *be a cyclic group,* $|C| = n$. *Let* V *be a finite symplectic vector space. Assume* C *acts on* V *preserving the symplectic form. Assume that* V *is an irreducible and faithful* C*-module. Set* $|V| = e^2$. *Then* n *divides* $e + 1$.

Proof. See [4], Satz 9.23. □

We say that a group G acts *Frobeniusly* on a vector space V if every nontrivial element of G acts fixed-point-free on V.

Lemma 2.2.3. *Let* V *be a finite symplectic vector space. Let* G *be a finite solvable group that acts faithfully, irreducibly and symplectically on* V. *Set* $|V| = e^2$. *Assume* $\mathbf{F}(G)$ *acts Frobeniusly on* V. *Let* M *be a normal cyclic subgroup of* $\mathbf{F}(G)$. *Then* $|M| \leq e + 1$.

Proof. Set $F = \mathbf{F}(G)$. Since F is normal in G, by Clifford theory we have that V decomposes as a direct sum of irreducible F-modules, each of the same size.

Our hypothesis is that F acts Frobeniusly on V. Assume that V is not an irreducible F-module. By Clifford theory we have that V decomposes as a direct sum of at least 2 irreducible F-modules, each of the same size. Since F acts Frobeniusly on V, we have that $|F| \leq |U|$, where U is one of the irreducible components of the F-module V. Thus $|F| \leq e$. Since M is a subgroup of F, it follows that $|M| \leq e < e + 1$.

We may assume that V is an irreducible F-module. By hypothesis, we have that V is a symplectic space and that G acts on V preserving the symplectic form. If V is not an irreducible M-module, then as before we can conclude that $|M| \leq e$. So we may assume that V is an irreducible M-module. Then by Lemma 2.2.2 we have that $|M|$ divides $e + 1$. □

Lemma 2.2.4. *Assume* G *is a finite solvable group. Then*

$$|G| \text{ divides } |\operatorname{Aut}(\mathbf{F}(G))||Z(\mathbf{F}(G))|.$$

Proof of Theorem 2.2.1. Since the hypothesis of Theorem 2.2.1 implies the hypothesis of Lemma 2.1.2, we will study the cases given by the conclusion of Lemma 2.1.2.

Claim 2.2.5. $e^2 - 1 = q^n - 1$ *divides the order of* G.

Proof. By Hypothesis, G acts transitively on $V^\#$ and $e^2 = |V|$. Therefore $e^2 - 1$ divides the order of G. □

Claim 2.2.6. *Assume that* G *is a subgroup of the semi-linear group* $\Gamma(V)$. *Also assume that* G *acts transitively on* $V^\#$. *Set* $H = G \cap U$, *where* U *is as in* (2.1.1). *Then* H *is a normal cyclic subgroup of* G *and* G/H *acts faithfully on* H.

Proof. Since U is a normal cyclic subgroup of $\Gamma(V)$, we have that $H = U \cap G$ is a normal cyclic subgroup of G. Since $\Gamma(V)/U$ is a cyclic group of order n, G/H is a cyclic group of order a divisor of n.

Since H is a cyclic normal subgroup of G, we have that G/H acts on H by conjugation. Suppose that G/H does not act faithfully on H. Let K/H be the kernel of the action and $t = |K/H|$. Thus G/K acts faithfully on H. Since H is cyclic, $|G/K|$ divides $\varphi(|H|)$, where φ is the Euler function. Since K/H is the kernel of the action of G/H on H, $t = |K/H|$ and G/H is isomorphic to a subgroup of $\mathcal{G}$, we have that H is isomorphic to a subgroup of the group of units $(\mathrm{GF}(q^{n/t}))^{\#}$ of a subfield of $\mathrm{GF}(q^n)$. Thus $|H|$ divides $q^{n/t} - 1$. Therefore we have that

$$|G| \text{ divides } (q^{n/t} - 1)t\varphi(q^{n/t} - 1). \tag{2.2.7}$$

Observe that $q^n - 1 = (q^{n/t} - 1)(q^{\frac{(t-1)n}{t}} + q^{\frac{(t-2)n}{t}} + \cdots + 1)$. Thus by (2.2.7) and Claim 2.2.5 we have that

$$(q^{\frac{(t-1)n}{t}} + q^{\frac{(t-2)n}{t}} + \cdots + 1) \text{ divides } t\varphi(q^{n/t} - 1). \tag{2.2.8}$$

Observe that $1 + q^{\frac{2n}{t}} > 2q^{n/t}$ since $(1 - q^{n/t})^2 > 0$. If $t > 2$, we have that

$$q^{\frac{(t-1)n}{t}} + q^{\frac{(t-2)n}{t}} + \cdots + 1 > tq^{n/t} > t\varphi(q^{n/t} - 1).$$

That can't hold by (2.2.8). Thus $t = 2$. If q is odd, then $q^{n/2} - 1$ is even. That implies

$$\varphi(q^{n/2} - 1) \leq \frac{1}{2}(q^{n/2} - 1).$$

But then

$$q^{n/2} + 1 > q^{n/2} - 1 \geq 2\varphi(q^{n/2} - 1).$$

By (2.2.8) we conclude that q must be even. Since $\varphi(q^{n/2} - 1) \leq q^{n/2} - 1$, by (2.2.8) we have that

$$q^{n/2} + 1 \bigm| 2\varphi(q^{n/2} - 1) \leq 2(q^{n/2} - 1).$$

Thus $q^{n/2} + 1 = 2\varphi(q^{n/2} - 1)$ and q can not be even. Therefore $t = 1$ and so the action of G/H on H is faithful. □

Case 2.2.9. *Assume (i) of Lemma 2.1.2, i.e., assume G is a subgroup of the semilinear group $\Gamma(V)$. Then*
(i) $e = 2$ and $G \cong \mathbf{Z}_3$ or $G \cong S_3$.
(ii) $e = 3$ and $G \cong \mathbf{Q}_8$.

Proof. By Claim 2.2.6, we have that $H = G \cap U$ is a normal cyclic subgroup of G and G/H acts faithfully on H.

Because H is cyclic, we have then $|\operatorname{Aut}(H)| = \varphi(|H|)$, where φ is the Euler function. Since G/H acts faithfully on H, we have that

$$|G| \leq |H|\varphi(|H|).$$

If $|H|$ is not a prime number and $|H| \neq 4$, then $\varphi(|H|) \leq |H| - 3$. By Lemma 2.2.3 we have that $|H| \leq e + 1$. So

$$\varphi(|H|) \leq |H| - 3 \leq e + 1 - 3 < e - 2.$$

But then $|G| \leq |H|\varphi(|H|) < e^2 - 1$, a contradiction with Claim 2.2.5. So we can assume that either $|H|$ is a prime number or $|H| = 4$.

Suppose that $|H|$ is a prime number q. Then by Lemma 2.2.3 we have that $q \leq e + 1$. If $q < e + 1$, then $q \leq e$ and

$$|G| \leq |H|\varphi(|H|) = q(q-1) \leq e(e-1) < e^2 - 1.$$

But that is a contradiction with Claim 2.2.5. Thus $q = e + 1$. Observe that $q(q-2) = e^2 - 1$. Because of Claim 2.2.5, we have that $q(q-2) = e^2 - 1$ divides $q(q-1)$. Since $e > 1$, $q = 3$ and $e = 2$. Thus $|H| = 3$ if $|H|$ is a prime number.

It remains to consider $|H| = 4$. By Lemma 2.2.4 $|G|$ divides $|H|\varphi(|H|) = 8$. By Claim 2.2.5, we have that $e^2 - 1$ divides $|G|$. Since $|G|$ divides 8, then $e^2 - 1 = |V^{\#}|$ is 2, 4 or 8. Since V is a symplectic vector space, $|V| = q^2$ for some power of a prime q. It follows that $|V^{\#}| = 8$ and $G \leq \mathrm{SL}(2,3)$. Thus $G \cong \mathbf{Q}_8$. □

Case 2.2.10. *Assume case (ii) of Lemma 2.1.2. Then*
(v) $q^n = 3^4$, $\mathbf{F}(G)$ *is extra-special of order* 2^5, $|\mathbf{F}_2(G)/\mathbf{F}(G)| = 5$ *and* $G/\mathbf{F}_2(G) \leq \mathbf{Z}_2$.

Proof. Since G acts symplectically and faithfully on a vector space V and $|V| = 3^4$, we have that G is isomorphic to a subgroup of $\mathrm{Sp}(3,4)$. Using the Atlas, we can check that $G/\mathbf{F}_2(G) \leq \mathbf{Z}_2$. □

Case 2.2.11. *Assume case (iii) of Lemma 2.1.2. Then* $e \in \{5,7\}$. *Furthermore one of the following holds*
(ii) $e = 3$ *and* $G \cong \mathrm{SL}(2,3)$.
(iii) $e = 5$ *and* $G \cong \mathrm{SL}(2,3)$.
(iv) $e = 7$ *and* $G \cong \widetilde{\mathrm{GL}}(2,3)$.

Proof. By Lemma 2.1.2 (iii), we have that $e^2 \in \{3^2, 5^2, 7^2, 11^2, 23^2\}$. Thus $G \leq \mathrm{Sp}(2,e)$. Recall that $\mathbf{F}(G) = \mathbf{Q}_8 T$, where $T \leq \mathbf{Z}(\mathrm{GL}(V))$. Thus $T = \mathbf{Z}(\mathrm{GL}(V)) \cap \mathrm{Sp}(2,e) = \mathbf{Z}(\mathrm{Sp}(2,e))$. Since $\mathrm{Sp}(2,e) = \mathrm{SL}(2,e)$ and $|\mathbf{Z}(\mathrm{SL}(2,e))| = 2$ by Lemma 2.1.3 (i) and (iii), we have that $|T| = 1$ or $|T| = 2$. Since $T \cap \mathbf{Q}_8 = \mathbf{Z}(\mathbf{Q}_8)$, it follows that $T = \mathbf{Z}(\mathbf{Q}_8)$. Thus $\mathbf{F}(G) = \mathbf{Q}_8$.

Since $\mathrm{Aut}(\mathbf{Q}_8) \cong S_4$ (see Exercise 5.3.4 of [7]), we have that $|\mathrm{Aut}(\mathbf{Q}_8)| = 24$. Since $\mathbf{Z}(\mathbf{Q}_8)$ is cyclic of order 2, by Claim 2.2.4 we have that $|G|$ is a divisor of 48. By Claim 2.2.5 we have that $e^2 - 1$ has to divide 48. Therefore $e \in \{3,5,7\}$.

By hypothesis G is isomorphic to a subgroup of $\mathrm{Sp}(V)$. Recall that $|V| = e^2$. By Lemma 2.1.3 we have that $\mathrm{SL}(2,e) = \mathrm{Sp}(2,e)$. Thus if $e = 3$ then $|G|$ divides $|\mathrm{SL}(2,3)| = 24$. Since G acts transitively on a set of 8 elements, we have that 8 divides $|G|$. Since $\mathbf{Q}_8/Z(\mathbf{Q}_8) \cong \mathbf{F}(G)/T$ is a faithful irreducible $G/\mathbf{F}(G)$-module, it follows that $G \cong \mathrm{SL}(2,3)$.

If $e = 5$, then G is isomorphic to a solvable subgroup of $\mathrm{SL}(2,5)$. By Claim 2.2.5 we have that the order of G is divisible by $24 = e^2 - 1$. Since $\mathrm{SL}(2,5)$ is not solvable (see Lemma 2.1.3 (iii)) and $|\mathrm{SL}(2,5)| = 5 \times 24$, we have that $|G| = 24$. Since $\mathbf{F}(G) \cong \mathbf{Q}_8$, and $|G| = 24$, we have that $G \cong \mathrm{SL}(2,3)$.

Similarly, if $e = 7$, G is isomorphic to a subgroup of $\mathrm{SL}(2,7)$. By Claim 2.2.5 we have that G is a solvable group of order divisible by $48 = 7^2 - 1$. Since $\mathrm{SL}(2,7)$ is not a solvable group (see Lemma 2.1.3 (iii)) and $|\mathrm{SL}(2,7)| = 7 \times 48$ we have that $|G| = 48$. By Lemma 2.1.6, we have that $G \cong \widetilde{\mathrm{GL}}(2,3)$. □

We have analyzed all the cases given in the conclusion of Lemma 2.1.2. So we have finished the proof. □

3. Theorem A

Our main result of this section is Theorem A. In Proposition 3.3.5, for each case (i) to (v) in the conclusion of Theorem A, we will give an example of a solvable group and a faithful character $\chi \in \mathrm{Irr}(G)$ satisfying the conditions in that case.

3.1. **General results.**

All the results in this section are proved in [1].

Lemma 3.1.1. *Assume that G is a finite group. Let L and N be normal subgroups of G such that L/N is an abelian chief factor of G. Let $\theta \in \mathrm{Irr}(L)$ be a G-invariant character such that θ_N is reducible. Then*

$$\theta\overline{\theta} = 1_N^L + \Phi$$

where Φ is either the zero function or a character of L and $[\Phi_N, 1_N] = 0$.

Proof. See Lemma 4.1 of [1]. ☐

Lemma 3.1.2. *Assume that G is a finite group and $\chi \in \mathrm{Irr}(G)$ is a faithful character. Let $\{\alpha_i \in \mathrm{Irr}(G)^\# \mid i = 1, \dots, n\}$ be the set of non-principal irreducible constituents of $\chi\overline{\chi}$. Then*

$$\mathbf{Z}(G) = \bigcap_{i=1}^{n} \mathrm{Ker}(\alpha_i).$$

Proof. See Lemma 5.1 of [1]. ☐

Lemma 3.1.3. *Assume that G is a finite solvable group and $\chi \in \mathrm{Irr}(G)$ with $\chi(1) > 1$. Let $\{\alpha_i \in \mathrm{Irr}(G)^\# \mid i = 1, \dots, n\}$ be the set of non-principal irreducible constituents of $\chi\overline{\chi}$. If $\mathrm{Ker}(\alpha_j)$ is not properly contained in $\mathrm{Ker}(\alpha_i)$ for all i, then $[\chi\overline{\chi}, \alpha_j] = 1$. Thus $1 \in \{[\chi\overline{\chi}, \alpha_i] \mid i = 1, \dots, n\}$.*

Proof. See Theorem C of [1]. ☐

3.2. **Proof of Theorem A.**

We will need two lemmas in the proof of Theorem A. These results are well known.

Lemma 3.2.1. *Let G be a solvable group with cyclic center $\mathbf{Z}(G)$. Assume that $E/\mathbf{Z}(G)$ is a chief factor of G. If $E/\mathbf{Z}(G)$ is a fully ramified section with respect to some $\Lambda \in \mathrm{Irr}(E)$, then $\mathbf{C}_G(E/\mathbf{Z}(G)) = E\mathbf{C}_G(E)$. Also $E/\mathbf{Z}(G)$ is a symplectic vector space and the conjugate action of G on $E/\mathbf{Z}(G)$ preserves the symplectic form.*

Lemma 3.2.2. *Let G be a group and H be a subgroup of G. If $\chi \in \mathrm{Irr}(G)$ and $\chi_H \in \mathrm{Irr}(H)$, then $\mathbf{C}_G(H) \leq \mathbf{Z}(\chi)$.*

Proof of Theorem A. Lemma 3.1.3 implies that $m = 1$.

Since χ is a faithful character, we have that $\mathbf{Z}(\chi) = \mathbf{Z}(G) = Z$. By Lemma 3.1.2 we have that $\mathrm{Ker}(\alpha) = Z$.

Claim 3.2.3. *Let N be a normal subgroup of G. Then $\chi_N \in \mathrm{Irr}(N)$ or $N \leq Z$.*

Proof. Assume N is not contained in Z. Since $\mathrm{Ker}(\alpha) = Z$ and N is normal in G, we have that $[\alpha_N, 1_N] = 0$. We conclude that $[(\chi\overline{\chi})_N, 1_N] = 1$. Thus χ_N is an irreducible character of N. ☐

Claim 3.2.4. *Let E/Z be a chief factor of G. Let $\lambda \in \mathrm{Irr}(Z)$ be such that $[\lambda, \chi_Z] \neq 0$. Then χ_E and λ are fully ramified with respect to E/Z. Therefore E/Z is a symplectic vector space and G acts on it preserving the symplectic form. Also $\chi(1)^2 = |E : Z|$.*

Proof. By Claim 3.2.3, we have that $\chi_E \in \mathrm{Irr}(E)$. By (1.0.1) we have that

$$\chi(1)^2 = 1 + m\lambda(1)$$

and therefore $\chi(1) > 1$. By hypothesis, G is solvable and so E/Z is abelian. Since $Z = \mathbf{Z}(G)$ and χ_Z reduces, by Theorem 6.18 of [5] we have that χ_E is fully ramified with respect to E/Z. Thus $\chi_Z = e\lambda$, where $e^2 = |E : Z|$. In particular, we have that $e = \chi(1)$. ☐

Claim 3.2.5. $(\chi\overline{\chi})_E = 1_Z^E$.

Proof. Since $\chi(1)^2 = |E : Z|$, by Lemma 3.1.1 we get the claim. ☐

Claim 3.2.6. $C_G(E/Z) = E$. *Thus G/E acts symplectically and faithfully on E/Z, transitively on $(E/Z)^{\#}$.*

Proof. By Lemma 3.2.1 we have that $\mathbf{C}_G(E/Z) = E\mathbf{C}_G(E)$. If $\mathbf{C}_G(E) \neq Z$, then Claim 3.2.3 implies that $\chi_{\mathbf{C}_G(E)} \in \mathrm{Irr}(\mathbf{C}_G(E))$. By Lemma 3.2.2 we have that

$$E \leq \mathbf{C}_G(\mathbf{C}_G(E)) \leq Z(\chi) = \mathbf{Z}(G) = Z < E.$$

We conclude that $\mathbf{C}_G(E) = Z$. Thus $E = \mathbf{C}_G(E/Z)$. Therefore G/E acts faithfully on E/Z.

By Claim 3.2.5 we have that $\alpha_E = 1_Z^E - 1_E$. Since E is normal in G and $\alpha \in \mathrm{Irr}(G)$, by Clifford theory we have that G acts transitively on the set

$$\mathrm{Irr}(E \bmod Z)^{\#} = \{\gamma \in \mathrm{Irr}(E) \mid \gamma \neq 1, \mathrm{Ker}(\gamma) \geq Z\}.$$

Therefore by Lemma 5.2 of [3], we have that G acts transitively on $(E/Z)^{\#}$. The rest follows from Claim 3.2.4 ☐

Claim 3.2.7. *Set $\bar{G} = G/E$. Then one of the following holds*

(i) $e = 2$ and either $G \cong S_3$ or $G \cong \mathbf{Z}_3$.

(ii) $e = 3$ and either $\bar{G} \cong \mathbf{Q}_8$ or $\bar{G} \cong \mathrm{SL}(2,3)$.

(iii) $e = 5$ and $\bar{G} \cong \mathrm{SL}(2,3)$.

(iv) $e = 7$ and $\bar{G} \cong \widetilde{\mathrm{GL}}(2,3)$.

(v) $e = 9$, $\mathbf{F}(\bar{G})$ is an extra-special group of order 2^5, namely the central product of the quaternion group $\mathbf{Q}_8$ and the dihedral group of order 8, $|\mathbf{F}_2(\bar{G})/\mathbf{F}(\bar{G})| = 5$ and $G/\mathbf{F}_2(\bar{G})$ is a subgroup of the cyclic group of order 2.

Proof. By Claim 3.2.6 we have the hypotheses of Theorem 2.2.1. The rest follows from Theorem 2.2.1. ☐

Since $e^2 = |E/Z|$ and $\chi(1) = e$, Theorem A follows from Claim 3.2.7. ☐

3.3. **Examples.** The following is a set of examples of solvable groups with irreducible characters that satisfy the hypotheses of Theorem A. We will use the notation of Theorem A in this subsection. We will start with some lemmas that we will use in the construction of some of the examples.

Lemma 3.3.1. *Let E be an extra-special group of order p^{2n+1} and exponent p, for some odd prime p. Observe that $E/\mathbf{Z}(E)$ is a vector space of dimension $2n$ over* $\mathrm{GF}(p)$ *with a symplectic form. Then the following short exact sequence*

$$0 \to \mathrm{Inn}(E) \to \mathrm{Aut}(E \bmod \mathbf{Z}(E)) \to \mathrm{Sp}(2n,p) \to 0$$

splits.

Proof. See [8] □

Lemma 3.3.2. *Let G be a finite group. Let $E/\mathbf{Z}(G)$ be an abelian chief factor of G. Assume that the action by conjugation of G/E on $(E/\mathbf{Z}(G))^{\#}$ is transitive. Let $\chi \in \mathrm{Irr}(G)$ be a faithful character of G. Assume that $\chi_E \in \mathrm{Irr}(E)$ and $\chi(1) > 1$. Then χ satisfies* (1.0.1), *i.e. $\eta(\chi) = 1$.*

Proof. Set $Z = \mathbf{Z}(G)$. Let $\lambda \in \mathrm{Irr}(Z)$ be a character of Z such that $[\chi_Z, \lambda] \neq 0$. Since $\chi_E \in \mathrm{Irr}(E)$ and $Z = \mathbf{Z}(G) \leq \mathbf{Z}(E)$, by Corollary 2.30 of [5] we have that $\chi(1)^2 \leq |E : Z|$. Observe that χ_Z is a reducible character since $\chi(1) > 1$ and $Z = \mathbf{Z}(G)$ is abelian. Observe also that $1_Z^E(1) = |E : Z| \geq \chi(1)^2$. Since $\chi(1)^2 \leq |E : Z|$, by Lemma 3.1.1 we conclude that

$$(\chi\overline{\chi})_E = 1_Z^E. \tag{3.3.3}$$

In particular, $\chi(1)^2 = |E : Z|$.

Let $\alpha \in \mathrm{Irr}(G)^{\#}$ be such that $[\alpha, \chi\overline{\chi}] \neq 0$. Observe such α exists since $\chi(1) > 1$. By Lemma 3.1.2, we have that $\mathrm{Ker}(\alpha) \geq Z$. By (3.3.3) we have that E is not contained in $\mathrm{Ker}(\alpha)$. Thus the irreducible constituents of α_E are elements of $\mathrm{Irr}(E \bmod Z)^{\#}$. By Clifford theory we have that the irreducible constituents of α_E are G-conjugate. Our hypothesis is that G/E acts transitively on $(E/Z)^{\#}$. Thus G/E acts transitively on the set $\mathrm{Irr}(E \bmod Z)^{\#}$. We conclude that the set of irreducible constituents of α_E is $\mathrm{Irr}(E \bmod Z)^{\#}$. Therefore $\alpha(1) \geq |E : Z| - 1$. Since $\chi(1)^2 = |E : Z|$, α is an irreducible constituent of $\chi\overline{\chi}$, $\alpha(1) \geq |E : Z| - 1$ and $[1_G, \chi\overline{\chi}] = 1$, we have that

$$\chi\overline{\chi} = 1_G + \alpha.$$

□

Lemma 3.3.4. *Assume the notation of Theorem A. Let $p \in \{3, 5, 7\}$. There exists a group G that satisfies the hypotheses of Theorem A and such that $\chi(1) = p$ and $|G : \mathbf{Z}(G)| = p^2 \times (p^2 - 1)$. In particular*

(i) If $p = 3$, then $\bar{G} \cong \mathbf{Q}_8$.
(ii) If $p = 5$, then $\bar{G} \cong \mathrm{SL}(2,3)$.
(iii) If $p = 7$, then $\bar{G} \cong \widetilde{\mathrm{GL}}(2,3)$.

Proof. Let E be an extra-special group of order p^3 and exponent p, where $p \in \{3, 5, 7\}$.

Observe that $\mathrm{SL}(2,3)$ has a subgroup of order 8, namely the quaternion group $\mathbf{Q}_8$. We can check that $\mathrm{SL}(2,5)$ has a subgroup isomorphic to $\mathrm{SL}(2,3)$ (see Lemma 2.1.3 (v)), and thus has a subgroup of order 24. Also $\mathrm{SL}(2,7)$ has a subgroup of order 48, namely $\widetilde{\mathrm{GL}}(2,3)$ (see Lemma 2.1.6). Thus for $p \in \{3, 5, 7\}$, there exists a solvable subgroup H of $\mathrm{SL}(2,p)$ of order $p^2 - 1$.

Set $V = E/\mathbf{Z}(E)$. Observe that V is a vector space of dimension 2 over $GF(p)$ and V has a symplectic form, namely $(\overline{v}, \overline{u}) \mapsto [v, u]$. Denote by $\mathrm{Sp}(2,p)$ the group of automorphisms of V that preserves the symplectic form of V. Since

$\mathrm{SL}(2,p) = \mathrm{Sp}(2,p)$ (see Lemma 2.1.3 (i)), we may assume that H is a subgroup of $\mathrm{Sp}(2,p)$. By Lemma 3.3.1 we may assume that $H \leq \mathrm{Aut}(E \bmod \mathbf{Z}(E))$. We can check that H acts transitively on $V^{\#}$.

Let $G = H \ltimes E$ be the semidirect product of E by H. Observe that $\mathbf{Z}(G) = \mathbf{Z}(E)$. Set $Z = \mathbf{Z}(G)$. Let $\Lambda \in \mathrm{Irr}(E)$ be a character of degree p. Observe that Λ is a G-invariant character. Since $(|H|, |E|) = 1$, and Λ is a G-invariant character, Λ extends to G (see Theorem 13.3 of [5]). Let $\chi \in \mathrm{Irr}(G)$ be an extension of Λ.

Observe that $\mathrm{Ker}(\chi)$ is a normal subgroup of G and acts on E by conjugation. Thus $\mathrm{Ker}(\chi)$ is normal in $\mathrm{Ker}(\chi)E$. Since E is also normal in $\mathrm{Ker}(\chi)E$, we have $\mathrm{Ker}(\chi) \leq \mathbf{C}_G(E) \leq E$. Since $\mathrm{Ker}(\chi) \cap E = \mathrm{Ker}(\Lambda) = 1$, it follows that χ is a faithful character.

Observe that we have the hypothesis of Lemma 3.3.2. Thus the result follows from that Lemma. □

Proposition 3.3.5. *For each of the cases (i) - (v) in the conclusion of Theorem A, there exists a solvable group and a faithful character $\chi \in \mathrm{Irr}(G)$ satisfying the conditions in that case.*

Proof. **(i)** Let $G = \mathrm{GL}(2,3)$ and $\chi \in \mathrm{Irr}(G)$ be a character of degree 2. We can check that χ satisfies (1.0.1). We can check that $G/\mathbf{Z}(G) \cong S_4$.

Let $G = \mathrm{SL}(2,3)$ and $\chi \in \mathrm{Irr}(G)$ be a character of degree 2. We can check that χ satisfies (1.0.1). We can check that $G/\mathbf{Z}(G) \cong A_4$.

(ii) Lemma 3.3.4 gives us a solvable group satisfying (ii) of Theorem A with $\bar{G} \cong \mathbf{Q}_8$. It remains to prove that there exist a group G and a faithful character χ satisfying (ii) such that $\bar{G} \cong \mathrm{SL}(2,3)$.

Let E be an extraspecial group of order 27 and exponent 3. Then $E/\mathbf{Z}(E) = V$ may be regard as a vector space of dimension 2 over GF(3), the finite field with 3 elements. Observe that $\mathrm{SL}(2,3)$ acts on V transitively. By Lemma 2.1.3 (i) we have that $\mathrm{SL}(2,3) = \mathrm{Sp}(2,3)$. Also by Lemma 2.1.3 (v), we have that $\mathrm{SL}(2,3) \cong \mathbf{Q}_8 \rtimes \mathbf{Z}_3$. We will assume that $\mathrm{SL}(2,3) = \mathbf{Q}_8 \rtimes \mathbf{Z}_3$ By Lemma 3.3.1 we may regard $\mathrm{SL}(2,3)$ as a subgroup of $\mathrm{Aut}(E \bmod \mathbf{Z}(E))$.

Let $G = \mathrm{SL}(2,3) \ltimes E$. Observe that E/Z is a chief factor of G since $\mathrm{SL}(2,3)$ acts transitively on $(E/Z)^{\#}$. Let $\Lambda \in \mathrm{Irr}(E)$ be a character of degree 3. Observe that Λ is G-invariant. Since $(|\mathbf{Q}_8|, |E|) = 1$ and Λ is G-invariant, we have that Λ extends to $\mathbf{Q}_8 \ltimes E$. Since $\mathbf{Z}_3$ is cyclic and Λ is G-invariant, Λ extends to $\mathbf{Z}_3 \ltimes E$. Thus Λ extends to G (see Corollary 11.31 of [5]). Let $\chi \in \mathrm{Irr}(G)$ be an extension of Λ. We can check that χ is a faithful character. By Lemma 3.3.2 it follows that $\eta(\chi) = 1$ and we are done with this case.

(iii) and (iv) This is Lemma 3.3.4 (ii) and (iii).

(v) Using the Atlas, we can check that $\mathrm{Sp}(4,3)$ has a subgroup H such that $\mathbf{F}(H)$ is an extra-special group of order 2^5, namely the central product of the quaternion group $\mathbf{Q}_8$ and the dihedral group of order 8, $|\mathbf{F}_2(H)/\mathbf{F}(H)| = 5$ and $H/\mathbf{F}_2(H)$ is a subgroup of the cyclic group of order 2.

Let E be an extraspecial group of order 3^5 and exponent 3, and let $\Lambda \in \mathrm{Irr}(E)$ be a character with degree 9. By Lemma 3.3.1, we may assume that H is a subgroup of $\mathrm{Aut}(E \bmod \mathbf{Z}(E))$. Set $G = H \ltimes E$. Observe that Λ is G-invariant. Since $(|H|, |E|) = 1$ and Λ is G-invariant, Λ extends to G. Let $\chi \in \mathrm{Irr}(G)$ be an extension of Λ. Lemma 3.3.2 implies that (1.0.1) holds for χ and so we are done. □

4. THEOREM B

In this section we study the solvable groups G that have an irreducible character χ such that $\chi\overline{\chi}$ has exactly 2 distinct non-principal irreducible constituents. Theorem B will be proved in Subsection 4.2. Examples of groups satisfying the hypotheses of Theorem B will be provided in Subsection 4.2. If, in addition, the group G is also nilpotent, the structure of the group is very constrained. We will classify those groups in Subsection 4.3

4.1. **Hypothesis 4.1.1.**

Hypothesis 4.1.1. *Let G be a finite solvable group. Let $\chi \in \mathrm{Irr}(G)$ be a faithful character. Assume*

$$\chi\overline{\chi} = 1_G + m_1\alpha_1 + m_2\alpha_2 \tag{4.1.2}$$

where $\alpha_1, \alpha_2 \in \mathrm{Irr}(G)$ are non-principal characters, m_1 and m_2 are strictly positive integers. Set $Z = \mathbf{Z}(G)$. Let $\lambda \in \mathrm{Irr}(Z)$ be the character such that $[\chi_Z, \lambda] \neq 0$.

4.2. **Derived length.**

Lemma 4.2.1. *Assume Hypothesis 4.1.1. Let N be a normal subgroup of G Then $\chi_N \in \mathrm{Irr}(N)$ if and only if $N \not\leq \mathrm{Ker}(\alpha_1)$ and $N \not\leq \mathrm{Ker}(\alpha_2)$.*

Proof. Observe that

$$[(\chi\overline{\chi})_N, 1_N] = [1_N + m_1(\alpha_1)_N + m_2(\alpha_2)N, 1_N] = 1 + m_1[(\alpha_1)_N, 1_N] + m_2[(\alpha_2)_N, 1_N].$$

Thus $[(\chi\overline{\chi})_N, 1_N] = [\chi_N, \chi_N] = 1$ if and only if $[(\alpha_1)_N, 1_N] = 0$ and $[(\alpha_2)_N, 1_N] = 0$. Therefore $[\chi_N, \chi_N] = 1$ if and only if $N \not\leq \mathrm{Ker}(\alpha_1)$ and $N \not\leq \mathrm{Ker}(\alpha_2)$. Since $[\chi_N, \chi_N] = 1$ if and only if $\chi_N \in \mathrm{Irr}(N)$, the result follows. □

Lemma 4.2.2. *Let T be a finite group and Θ a character of T. Suppose that $T = R \times S$. Assume that $\Theta(g) = 0$ if $g \in T \setminus R$ or if $g \in T \setminus S$. Then*

$$\Theta(g) = \begin{cases} 0 & \text{if } g \neq 1, \\ \Theta(1) & \text{otherwise.} \end{cases} \tag{4.2.3}$$

Thus Θ is a multiple of the regular character.

Proof. Since $\Theta(g) = 0$ for $g \in T \setminus R$ and $R \cap S = 1$, $\Theta(s) = 0$ for all $s \in S \setminus \{1\}$. Also $\Theta(t) = 0$ for all $t \in T \setminus S$. Thus (4.2.3) holds. □

Lemma 4.2.4. *Assume Hypothesis 4.1.1. Then either $\mathrm{Ker}(\alpha_1) = Z$ or $\mathrm{Ker}(\alpha_2) = Z$.*

Proof. Assume that $\mathrm{Ker}(\alpha_1) \neq Z$ and $\mathrm{Ker}(\alpha_2) \neq Z$.

Let R/Z be a chief factor of G, where $R \leq \mathrm{Ker}(\alpha_1)$. Note that such an R exists since $Z = \mathrm{Ker}(\alpha_1) \cap \mathrm{Ker}(\alpha_2)$ by Lemma 3.1.2, and $\mathrm{Ker}(\alpha_1) \neq Z$ by assumption. Observe also that $R \not\leq \mathrm{Ker}(\alpha_2)$. Similarly, let S/Z be a chief factor of G such that $S \leq \mathrm{Ker}(\alpha_2)$. Set $T = RS$. Since $\mathrm{Ker}(\alpha_1) \cap \mathrm{Ker}(\alpha_2) = Z$, we have that $R \cap S = Z$. Hence $T \not\leq \mathrm{Ker}(\alpha_1)$ and $T \not\leq \mathrm{Ker}(\alpha_2)$. By Lemma 4.2.1, we have that $\chi_T \in \mathrm{Irr}(T)$.

Since S/Z is a chief factor and $T = RS$ with $R \cap S = Z$, we have that T/R is a chief factor of G. Let $\psi \in \mathrm{Irr}(R)$ be such that $[\chi_R, \psi] \neq 0$. By Theorem 6.18 of [5] one of the following holds

a) $\chi_R = \psi$.

b) $\chi_R = e_1\psi$ for some $\psi \in \mathrm{Irr}(R)$ and some integer $e_1 > 0$ such that ${e_1}^2 = |T : R|$.

c) $\psi^T = \chi_T$.

Observe that $\chi_R = \psi$ can not hold since $R \leq \mathrm{Ker}(\alpha_1)$ and Lemma 4.2.1. Observe also that b) and c) imply that $\chi(g) = 0$ if $g \in T \setminus R$. Similarly, we can check that $\chi(g) = 0$ if $g \in T \setminus S$. Since R/Z, S/Z are chief factors of G, they are elementary abelian. Thus we have

$$\chi\overline{\chi}(e) = \begin{cases} 0 & \text{if } e \in T \setminus R \\ 0 & \text{if } e \in T \setminus S \\ \chi(1)^2 & \text{if } e \in R \cap S = Z. \end{cases}$$

Recall that R and S are normal subgroups of G with $R \cap S = Z$ and $T = RS$. Then we have that $T/Z = R/Z \times S/Z$. By Lemma 4.2.2 we have that $\chi\overline{\chi}$ is a multiple of the regular character of $RS \bmod Z$, i.e. the regular character of RS/Z inflated to RS. Thus $\chi(1)^2 \geq |T : Z|$. Since $\chi_T \in \mathrm{Irr}(T)$ and $Z = \mathbf{Z}(G)$, we have that $\chi(1)^2 \leq |T : Z|$. Therefore $(\chi\overline{\chi})_T = 1_Z^T$ is the regular character of $T \bmod Z$. Because all the characters of $\mathrm{Irr}(T \bmod Z)$ are linear, and thus appear with multiplicity 1 in 1_Z^T, and $(\chi\overline{\chi})_T = 1_Z^T$, we have that $m_1 = m_2 = 1$.

Observe that

$$\begin{aligned}[(\chi\overline{\chi})_R, 1_R] &= [((\chi\overline{\chi})_T)_R, 1_R] \\ &= [(1_Z^T)_R, 1_R] = |T : R|.\end{aligned}$$

Recall that $R \leq \mathrm{Ker}(\alpha_1)$ and $R \not\leq \mathrm{Ker}(\alpha_2)$. Applying (4.1.2) we get that

$$1 + \alpha_1(1) = [(\chi\overline{\chi})_R, 1_R] = |T : R|.$$

Similarly we can check that

$$1 + \alpha_2(1) = |T : S|.$$

Thus

$$\chi(1)^2 = |T : Z| = 1 + \alpha_1(1) + \alpha_2(1) = |T : R| + |T : S| - 1.$$

Because $T = RS$ and $R \cap S = Z$, we have that $|T : S| = |R : Z|$. Observe that

$$|T : Z| = |T : R||R : Z|.$$

So we get

$$|T : R||R : Z| = |T : R| + |R : Z| - 1.$$

Therefore

$$(|T : R| - 1)(|R : Z| - 1) = 0.$$

This is impossible since neither $|T : R|$ nor $|R : Z|$ is 1. Thus either $\mathrm{Ker}(\alpha_1) = Z$ or $\mathrm{Ker}(\alpha_2) = Z$. □

We are going to describe how G looks, depending on $\mathrm{Ker}(\alpha_1)$ and $\mathrm{Ker}(\alpha_2)$.

Proposition 4.2.5. *Assume Hypothesis 4.1.1. Assume also that*

$$\mathrm{Ker}(\alpha_1) = \mathrm{Ker}(\alpha_2) = Z.$$

Let E/Z be a chief factor of G. Then E/Z is a fully ramified section with respect to χ_E and λ. Furthermore G/E acts faithfully and symplectically on E/Z, with 2 orbits on $(E/Z)^\#$. Thus $\mathrm{dl}(G) \leq 18$. Also $m_1 = m_2 = 1$, $\chi(1)^2 = |E : Z|$ and $\chi(1)$ is a power of a prime.

Proof. By Lemma 4.2.1 we have that $\chi_E \in \mathrm{Irr}(E)$. Since $\chi_E(1) \neq 1$, E/Z is a fully ramified section with respect to χ_E and λ. By Lemma 3.2.1 we have that $\mathbf{C}_G(E/\mathbf{Z}(G)) = E\mathbf{C}_G(E)$ and E/Z has the structure of a symplectic vector space. Since χ is a faithful character, we have that $\mathbf{Z}(\chi) = \mathbf{Z}(G) = Z$. By Lemma 3.2.2 we have that $\mathbf{C}_G(E) \leq \mathbf{Z}(\chi) = Z$. Thus $\mathbf{C}_G(E/Z) = E$. We conclude that G/E acts faithfully and symplectically on E/Z.

Since E/Z is a fully ramified section with respect to χ_E and λ, we have that

$$(\chi\overline{\chi})_E = 1_E^Z. \tag{4.2.6}$$

In particular, $\chi(1)^2 = |E : Z|$. Since E/Z is a chief factor of a solvable group, $|E : Z|$ is a power of a prime number. Therefore $\chi(1)$ is a power of a prime number.

By (4.1.2) and (4.2.6) we have that

$$1_E^Z = (\chi\overline{\chi})_E = (1_G + m_1\alpha_1 + m_2\alpha_2)_E = 1_E + m_1(\alpha_1)_E + m_2(\alpha_2)_E. \tag{4.2.7}$$

By Clifford Theory we have that the irreducible constituents of $(\alpha_1)_E$ form a G-orbit. Similarly, the irreducible constituents of $(\alpha_2)_E$ form a G-orbit. If the irreducible constituents of $(\alpha_1)_E$ are the irreducible constituents of $(\alpha_2)_E$, then by (4.2.7) G/E acts transitively on $1_E^Z \setminus 1_E$. Since χ_E is irreducible, by Lemma 3.3.2 and Theorem A it follows that $\chi\overline{\chi} = 1_G + \alpha$, where $\alpha \in \mathrm{Irr}(G)$. Thus the irreducible constituents of $(\alpha_1)_E$ and $(\alpha_2)_E$ form two distinct G-orbits. By (4.2.7), we have that the number of orbits of G on $\mathrm{Irr}(E \bmod Z)^{\#}$ is 2. Therefore the number of orbits of G on $(E/Z)^{\#}$ is 2.

Since G/E acts faithfully and irreducibly on E/Z, with 2 orbits on $(E/Z)^{\#}$, by Theorem 5.4 of [6] we have that $\mathrm{dl}(G/E) \leq 16$. Therefore $\mathrm{dl}(G) \leq 16 + 2 = 18$. Observe that $m_1 = m_2 = 1$ by Lemma 3.1.3. ☐

Remark. We will give an example of a group satisfying the hypotheses of Proposition 4.2.5 in Proposition 4.3.3.

Proposition 4.2.8. *Assume Hypothesis 4.1.1. Then the order of G is even.*

Proof. Let $\lambda \in \mathrm{Irr}(Z)$ be such that $[\chi_Z, \lambda] \neq 0$. Let E/Z be a chief factor of G.

If $2 \mid \chi(1)$, then $|G|$ has to be even since $\chi(1) \mid |G|$. So we may assume that $\chi(1)$ is an odd number.

If α_i is a real character, for some i, then G contains a conjugacy class with elements of order 2. Thus G has to be of even order. We may assume that α_1 and α_2 are not real characters.

Since $\chi\overline{\chi}$ is a real character and α_1 and α_2 are not real characters, by (4.1.2) we conclude that $\overline{\alpha_1} = \alpha_2$. Thus $\mathrm{Ker}(\alpha_1) = \mathrm{Ker}(\alpha_2)$ and $\alpha_1(1) = \alpha_2(1)$. Since $Z = \mathrm{Ker}(\alpha_1) \cap \mathrm{Ker}(\alpha_2)$ by Lemma 3.1.2, we conclude that $\mathrm{Ker}(\alpha_1) = \mathrm{Ker}(\alpha_2) = Z$. Thus by the previous Lemma, $\chi(1)$ is a power of an odd prime and $m_1 = m_2 = 1$. In particular 4 divides $\chi(1)^2 - 1$.

By (4.1.2) we have that

$$\chi(1)^2 = 1 + 2\alpha_1(1).$$

Thus $2\alpha_1(1) = \chi(1)^2 - 1$. Since 4 divides $\chi(1)^2 - 1$, we must have that 2 divides $\alpha_1(1)$. By Hypothesis 4.1.1 we have that $\alpha_1 \in \mathrm{Irr}(G)$. Thus 2 divides the order of G. ☐

Proposition 4.2.9. *Assume Hypothesis 4.1.1. Assume also that $K = \mathrm{Ker}(\alpha_1)$ is an abelian group properly containing Z. Let $\theta \in \mathrm{Irr}(K)$ be such that $[\chi_K, \theta] \neq 0$.*

Denote by $G_\theta = \{g \in G | \theta^g = \theta\}$ *the stabilizer in* G *of* θ. *Let* L/K *be a chief factor of* G. *Then*

(i) $\mathrm{Ker}(\alpha_2) = Z$.

(ii) $G = G_\theta L$ *and* $G_\theta \cap L = K$.

(iii) G_θ/K *acts faithfully and irreducibly on* L/K, *and transitively on* $(L/K)^\#$.

In particular, $\mathrm{dl}(G) \leq 6$ *and* $\chi(1)$ *is a power of a prime. Also* $m_1 = 1$.

Proof. Since $\mathrm{Ker}(\alpha_1) \neq Z$, by Lemma 4.2.4 it follows that $\mathrm{Ker}(\alpha_2) = Z$.

By Lemma 4.2.1 we have that $\chi_L \in \mathrm{Irr}(L)$. Since χ_K is reducible, K is abelian and $\mathbf{Z}(\chi) = Z \neq K$, by Theorem 6.18 of [5] we have that $\chi_L = \theta^L$ and that $G_\theta < G$. By Clifford theory it follows that χ is induced from an irreducible character of G_θ. By Exercise 5.7 of [5], we have that $G = G_\theta L$. Since L/K is an abelian group, $G_\theta \geq K$ and $G = G_\theta L$, we have that $G_\theta \cap L$ is a normal subgroup of G containing K. Observe that χ_{G_θ} is reducible since χ is induced from some character of G_θ and $G_\theta < G$. Thus $\chi_{G_\theta \cap L}$ is a reducible character, since $G_\theta \cap L \leq G_\theta$. Because $K \leq G_\theta \cap L \leq L$, $G_\theta \cap L$ is a normal subgroup of G and L/K is a chief factor of G, we have that either $G_\theta \cap L = L$ or $G_\theta \cap L = K$. If $G_\theta \cap L = L$ then $G = G_\theta L = G_\theta < G$. Thus $G_\theta \cap L = K$.

Observe that $\mathbf{C}_G(L/K)$ is a normal subgroup of G that contains L. Set $C = \mathbf{C}_G(L/K)$. Since $G = G_\theta L$, we have that $C = (C \cap G_\theta)L$. Observe that $C \cap G_\theta \trianglelefteq G_\theta$ and $[C \cap G_\theta, L] \leq [C, L] \leq K$. Thus $C \cap G_\theta \leq G_\theta$ is a normal subgroup of G. If $C \neq L$ then $C \cap G_\theta \leq G_\theta$ is a normal subgroup of G that properly contains K. Thus $\chi_{C \cap G_\theta} \in \mathrm{Irr}(C \cap G_\theta)$ by Lemma 4.2.1. This can not be because χ is induced from some character of G_θ and $C \cap G_\theta \leq G_\theta < G$. Thus $C \cap G_\theta = K$ and $C = L$. Therefore G_θ/K acts faithfully on L/K. Also G_θ/K acts irreducibly on L/K since $G = G_\theta L$ acts irreducibly on the chief factor L/K, $L = C = \mathbf{C}_G(L/K)$ and $C \cap G_\theta = K$.

By Lemma 3.1.1 we have that

$$(\chi\overline{\chi})_L = 1_K^L + \Phi, \tag{4.2.10}$$

where Φ is either the zero function or a character of L, and $[\Phi_K, 1_K] = 0$. Also, by (4.1.2) we have that

$$(\chi\overline{\chi})_L = 1_L + (m_1\alpha_1)_L + (m_2\alpha_2)_L. \tag{4.2.11}$$

Since $\mathrm{Ker}(\alpha_2) = Z < \mathrm{Ker}(\alpha_1)$, we have that $[(\alpha_2)_K, 1_K] = 0$. Thus by (4.2.10) and (4.2.11) it follows that the irreducible constituents of $(\alpha_1)_L$ form the set $\mathrm{Irr}(L \bmod K)^\#$. It follows that G acts transitively on $\mathrm{Irr}(L \bmod K)^\#$. Therefore G acts transitively on $(L/K)^\#$.

Since $\mathbf{C}_G(L/K) = L$, we have that that G/L acts faithfully on L/K and transitively on $(L/K)^\#$. Using Lemma 2.1.2, we can check that $\mathrm{dl}(G/L) \leq 4$. Since L/K is a chief factor of G and K is abelian, it follows that $\mathrm{dl}(L) \leq 2$. We conclude that $\mathrm{dl}(G) \leq 6$.

Since $\chi_L = \theta^L$ and $\theta(1) = 1$, we have that $\chi(1) = |L : K|$. Observe that $|L : K|$ is a power of a prime since L/K is a chief factor of a solvable group. Thus $\chi(1)$ is a power of a prime number. By Lemma 3.1.3 we have that $m_1 = 1$. □

Remark. In Proposition 4.3.4, we will prove that for any prime p and any nonzero positive integer m, there exist a group G and a character $\chi \in \mathrm{Irr}(G)$ with $\chi(1) = p^m$ satisfying the hypotheses of Proposition 4.2.9.

Lemma 4.2.12. *Assume that* G *is a group such that* $\mathbf{F}(G)$ *is an extra-special group of order* 2^5, $|\mathbf{F}_2(G)/\mathbf{F}(G)| = 5$ *and* $G/\mathbf{F}_2(G)$ *is a subgroup of the cyclic group of order 2.*

Then $\mathrm{cd}(G) \subseteq \{1, 2, 4, 5, 8, 10\}$. *Also, if* $\alpha \in \mathrm{Irr}(G)$ *and* $\alpha(1) = 8$, *then* $\mathrm{Ker}(\alpha) = 1$.

Proof. Observe that $\mathbf{Z}(G) = \mathbf{Z}(\mathbf{F}(G))$. Since $\mathbf{F}(G)$ is an extra-special group of order 2^5, there exists a unique $\Lambda \in \mathrm{Irr}(\mathbf{F}(G))$ such that $\Lambda(1) = 4$. This Λ is a G-invariant character. Since the p-Sylow subgroups of $G/\mathbf{F}(G)$ are cyclic for all primes p, and Λ is a G-invariant character, we have that Λ extends to G. Observe that $\mathrm{cd}(G/\mathbf{F}(G)) \subseteq \{1, 2\}$. Thus the degrees of the characters in $\mathrm{Irr}(G|\Lambda)$ lie in the set $\{4, 8\}$, while those of $\mathrm{Irr}(G \bmod \mathbf{F}(G))$ lie in $\{1, 2\}$.

Let $\lambda \in \mathrm{Lin}(\mathbf{F}(G) \bmod \mathbf{Z}(G))^{\#}$ be a linear character of $\mathbf{F}(G)$. Since $\mathbf{F}_2(G)/\mathbf{F}(G)$ acts without fixed points on $(\mathbf{F}(G)/\mathbf{Z}(\mathbf{F}(G)))^{\#}$, i.e on $\mathrm{Lin}(\mathbf{F}(G))^{\#}$, we have that λ induces irreducibly to $\mathbf{F}_2(G)$. If $\chi \in \mathrm{Irr}(G)$ lies above λ, then $\chi_{\mathbf{F}(G)}$ is the sum of $\chi(1)$ distinct linear characters of $\mathbf{F}(G) \bmod \mathbf{Z}(G)$. Because of $|G : \mathbf{F}(G)| \in \{5, 10\}$, we have that $\chi(1) \in \{5, 10\}$. □

Proposition 4.2.13. *Assume Hypothesis 4.1.1. Assume also that* $\mathrm{Ker}(\alpha_1)$ *is not abelian. Let* E/Z *be a chief factor of* G *such that* $E \leq \mathrm{Ker}(\alpha_1)$. *Let* F_2/E *be the Fitting subgroup of* G/E. *Then* $m_1 = 1$, $m_2 = 4$, $\alpha_1(1) = 3$ *and* F_2/E *is isomorphic to* Q_8. *Also, one of the following holds*

(i) $\chi(1) = 10$, $G/E \cong \mathrm{SL}(2, 3)$.

(ii) $\chi(1) = 14$, $G/E \cong \widetilde{\mathrm{GL}}(2, 3)$.

Proof. Set $K = \mathrm{Ker}(\alpha_1)$. Let $\theta \in \mathrm{Irr}(K)$ be a character such that $[\chi_K, \theta] \neq 0$. Since K is nonabelian and χ is faithful, χ_K is a sum of G-conjugate nonlinear characters, one of which is θ. Therefore $\theta(1) > 1$.

Let $\lambda \in \mathrm{Irr}(Z)$ be such that $[\lambda, \theta_Z] \neq 0$. By Lemma 4.2.4 we have that $\mathrm{Ker}(\alpha_2) = Z$.

Claim 4.2.14. *For any normal subgroup* N *of* G *such that* $Z < N \leq K$, *we have* $\theta_N \in \mathrm{Irr}(E)$. *In particular*

$$\Lambda = \theta_E \in \mathrm{Irr}(E). \tag{4.2.15}$$

Proof. Suppose that θ_N is a reducible character. Then $[\theta_N, \theta_N] > 1$. Since $[\chi_K, \theta] \neq 0$, we have that

$$[(\chi\overline{\chi})_N, 1_N] = [\chi_N, \chi_N] > [\chi_K, \chi_K] = [(\chi\overline{\chi})_K, 1_K].$$

Thus there exists an irreducible constituent α of $\chi\overline{\chi}$ such that $N \leq \mathrm{Ker}(\alpha)$ and $K \not\leq \mathrm{Ker}(\alpha)$. But the irreducible constituents of $\chi\overline{\chi}$ are 1_G, α_1 and α_2 and their kernels are G, K and Z respectively. We conclude that θ_N is an irreducible character. □

Since θ_E is not a linear character, E/Z is a chief factor of G and $Z = \mathbf{Z}(G)$, the section E/Z is fully ramified with respect to Λ and λ. Thus

$$\Lambda \text{ is } G\text{-invariant and } \Lambda(1)^2 = |E : Z|. \tag{4.2.16}$$

Claim 4.2.17. $\mathbf{C}_G(E/Z) = E$. *Thus* G/E *acts faithfully on the symplectic vector space* E/Z. *Also* G/E *acts transitively on* $(\mathrm{Lin}(E/Z))^{\#}$, *hence on* $(E/Z)^{\#}$.

Proof. We will first prove that $\mathbf{C}_G(E) = Z$. Observe that $C_G(E)$ is a normal subgroup of G. Suppose that $\mathbf{C}_G(E) \neq Z$. If $\mathbf{C}_G(E) \leq K$, by Claim 4.2.14 we have that $\theta_{\mathbf{C}_G(E)} \in \mathrm{Irr}(C_G(E))$. By Lemma 3.2.2 we have that

$$E \leq \mathbf{C}_K(\mathbf{C}_G(E)) \leq \mathbf{Z}(\theta).$$

Since $\Lambda = \theta_E \in \mathrm{Irr}(E)$ and $E \leq \mathbf{Z}(\theta)$, it follows that Λ is a linear character. Therefore $\theta(1) = 1$. But $\theta(1) > 1$. So we may assume that $\mathbf{C}_G(E) \not\leq K$. By Lemma 4.2.1 we have that $\chi_{\mathbf{C}_G(E)} \in \mathrm{Irr}(\mathbf{C}_G(E))$. As before, by Lemma 3.2.2 we have that

$$E \leq \mathbf{C}_G(\mathbf{C}_G(E)) \leq \mathbf{Z}(\chi) = Z < E,$$

a contradiction. Thus $\mathbf{C}_G(E) = Z$. By Lemma 3.2.1 we have that $\mathbf{C}_G(E/Z) = \mathbf{C}_G(E)E$. Since $\mathbf{C}_G(E) = Z$, it follows that $\mathbf{C}_G(E/Z) = E$.

By (4.2.15) and (4.2.16) we have that E/Z is a fully ramified section with respect to Λ and λ. Since $Z = \mathbf{Z}(G)$, we have

$$(\theta\overline{\theta})_E = \Lambda\overline{\Lambda} = 1_Z^E. \tag{4.2.18}$$

Since $[\Lambda, (\chi)_E] \neq 0$, we have $(\chi\overline{\chi})_E = 1_Z^E + \Phi$, where Φ is character of E. Since $K = \mathrm{Ker}(\alpha_1) \geq E$ and (4.1.2) holds we have

$$\begin{aligned}(\chi\overline{\chi})_E &= (1_G + m_1\alpha_1 + m_2\alpha_2)_E\\ &= (1 + m_1\alpha_1(1))1_E + m_2(\alpha_2)_E\\ &= 1_Z^E + \Phi.\end{aligned}$$

By Clifford Theory, the irreducible constituents of $(\alpha_2)_E$ are G-conjugates. Thus $(\mathrm{Irr}(E \bmod Z))^{\#}$ is a G-orbit. □

Claim 4.2.19. *Set* $e^2 = |E : Z|$ *and* $\bar{G} = G/E$. *Then one of the following holds:*

(i) $e = 2$, *and either* $\bar{G} \cong S_3$ *or* $\bar{G} \cong \mathbf{Z}_3$.

(ii) $e = 3$, *and either* $\bar{G} = \mathbf{Q}_8$ *or* $\bar{G} \cong \mathrm{SL}(2,3)$.

(iii) $e = 5$ *and* $\bar{G} \cong \mathrm{SL}(2,3)$.

(iv) $e = 7$ *and* $\bar{G} \cong \widetilde{\mathrm{GL}}(2,3)$.

(v) $e = 9$, $\mathbf{F}(\bar{G})$ *is an extra-special group of order* 2^5, $|\mathbf{F}_2(\bar{G})/\mathbf{F}(\bar{G})| = 5$ *and* $\bar{G}/\mathbf{F}_2(\bar{G})$ *is a subgroup of the cyclic group of order 2.*

Proof. This follows from Claim 4.2.17 and Theorem 2.2.1. □

Claim 4.2.20. *Let* $e \in \{2,3,5,7\}$. *Set* $\Lambda = (\theta)_E$. *Then* $\Lambda \in \mathrm{Irr}(E)$, $\Lambda(1) = e$ *and* Λ *extends to* G, *i.e. there exists* $\Lambda^e \in \mathrm{Irr}(G)$ *such that* $(\Lambda^e)_E = \Lambda$.

Proof. By (4.2.15) we have that $\Lambda = \theta_E \in \mathrm{Irr}(E)$. By (4.2.18) we have that $\Lambda(1) = e$.

Assume $e = 2$. Since the p-Sylow subgroups of G/E are cyclic for all primes p, by Corollary 11.22 and Corollary 11.31 of [5] we have that Λ extends to G.

Assume now that $e \in \{3,5,7\}$. Observe that for primes $p \neq 2$, the p-Sylow subgroups of G/E are cyclic. If we can prove that Λ extends to ET, where T is a 2-Sylow subgroup of G, then Λ would extend to G (see Corollary 11.31 of [5]). In our situation we have that either $TE/E \cong \mathbf{Q}_8$ or $TE/E \cong \mathbf{Q}_{16}$. It follows from the example on page 301 of [2] that $H^2(\mathbf{Q}_8, \mathbf{C}) = 1$ and $H^2(\mathbf{Q}_{16}, \mathbf{C}) = 1$. By Proposition 11.45 of [2] this implies that Λ extends to ET. Thus Λ extends to G. □

Claim 4.2.21. *Let* $e \in \{2,3,5,7\}$. *Let* $\Lambda^e \in \mathrm{Irr}(G)$ *be an extension of* $\Lambda \in \mathrm{Irr}(E)$. *Then*

$$\chi = \Lambda^e \psi \tag{4.2.22}$$

for some $\psi \in \mathrm{Irr}(G \bmod E)$. *Also*

(i) $\Lambda^e \overline{\Lambda^e} = 1_G + \alpha_2$.

(ii) $\psi\overline{\psi} = 1_G + \alpha_1$.

(iii) $\alpha_1\alpha_2 = \alpha_1(1)\alpha_2$ *and* $m_2 = 1 + \alpha_1(1)$.

Proof. Since $[\Lambda, \chi_E] \neq 0$ and Λ extends to $\Lambda^e \in \mathrm{Irr}(G)$, by Corollary 6.17 of [5] there exists $\psi \in \mathrm{Irr}(G \bmod E)$ such that (4.2.22) holds.

Observe that $\mathrm{Ker}(\Lambda^e) \leq E$ since Λ^e is an extension of Λ and G/E acts faithfully on E/Z (see Claim 4.2.17). Since χ is faithful and $\mathrm{Ker}(\Lambda^e) \leq E$, by (4.2.22) we have that $\mathrm{Ker}(\Lambda^e) = \mathrm{Ker}(\chi) \cap E = 1$. Thus Λ^e is a faithful character of G, E/Z is a chief factor of G, where $Z = \mathbf{Z}(G)$, and G/E acts on $(E/Z)^{\#}$ transitively (see Claim 4.2.17). By Lemma 3.3.2, we conclude that

$$\Lambda^e \overline{\Lambda^e} = 1_G + c_1\gamma$$

for some $\gamma \in \mathrm{Irr}(G)$ and some integer c_1. By Theorem A we have that $c_1 = 1$. By Lemma 3.1.2, we observe that

$$\mathrm{Ker}(\gamma) = Z. \tag{4.2.23}$$

Let $\psi\overline{\psi} = 1_G + \sum_{i=1}^{\eta(\psi)} d_i\delta_i$, for some integers $d_i > 0$ and some distinct characters $\delta_i \in \mathrm{Irr}(G)$ for $i = 1, \ldots, \eta(\psi)$. Since $\psi \in \mathrm{Irr}(G \bmod E)$, we have that $\delta_i \in \mathrm{Irr}(G \bmod E)$ for all i. Observe

$$\begin{aligned}
\chi\overline{\chi} &= \Lambda^e\psi\overline{\Lambda^e\psi} \\
&= (\Lambda^e\overline{\Lambda^e})(\psi\overline{\psi}) \\
&= (1_G + \gamma)(1_G + \sum_{i=1}^{\eta(\psi)} d_i\delta_i) \\
&= 1_G + \gamma + \sum_{i=1}^{\eta(\psi)} d_i\delta_i + \sum_{i=1}^{\eta(\psi)} d_i\gamma\delta_i \\
&= 1_G + m_1\alpha_1 + m_2\alpha_2
\end{aligned}$$

where the last equality is (4.1.2). Since $\mathrm{Ker}(\delta_1) \geq E$ and $\mathrm{Ker}(\gamma) = Z \not\leq E$ by (4.2.23), we have that $\gamma \neq \delta_1$. It follows that $\eta(\psi) = 1$.

Since $\mathrm{Ker}(\delta_1) \geq E$ and $\mathrm{Ker}(\gamma) = Z < E$, we have that $\mathrm{Ker}(\gamma\delta_1) \not\geq E$. Since $m_1 = 1$, $\mathrm{Ker}(\alpha_1) \geq E$ and $\mathrm{Ker}(\alpha_2) = Z$, it follows that $\delta_1 = \alpha_1$, $\gamma = \alpha_2$ and $\gamma\delta$ is a multiple of α_2. Thus $\gamma\delta = \alpha_1(1)\alpha_2$. Therefore $m_2 = 1 + \alpha_1(1)$. ☐

Claim 4.2.24. *Let* $e \in \{2,3,5,7\}$. *Set* $H = \mathrm{Ker}(\psi)$, *where* $\psi \in \mathrm{Irr}(G \bmod E)$ *is as in Claim 4.2.21. Let* $Z_H/H = \mathbf{Z}(G/H)$. *Then there exists* $L \trianglelefteq G$ *such that* L/Z_H *is a chief factor of* G. *Also* $E < L$, $\psi(1)^2 = |L : Z_H|$ *and one of the following holds*

(i) $\psi(1) = 2$ *and either* $G/L \cong \mathbf{Z}_3$ *or* $G/L \cong S_3$.

(ii) $\psi(1) = 3$ *and either* $G/L \cong \mathbf{Q}_8$ *or* $G/L \cong \mathrm{SL}(2,3)$.

(iii) $\psi(1) = 5$ *and* $G/L \cong \mathrm{SL}(2,3)$.

(iv) $\psi(1) = 7$ *and* $G/L \cong \widetilde{\mathrm{GL}}(2,3)$.

(v) $\psi(1) = 9$, $\mathbf{F}(G/L)$ *is an extra-special group of order* 2^5, $|\mathbf{F}_2(G/L)/\mathbf{F}(G/L)| = 5$ *and* $(G/E)/\mathbf{F}_2(G/L)$ *is a subgroup of the cyclic group of order 2.*

Proof. By Claim 4.2.21 (ii), we have that $\psi(1)^2 = 1 + \alpha(1) > 1$. Thus $\psi(1) > 1$. Since ψ is a faithful irreducible character of G/H and $\psi(1) > 1$, it follows that G/H is not abelian. Therefore there exists a normal subgroup L of G such that L/Z_H is a chief factor of G.

Observe that ψ is a faithful and irreducible character of $G/\operatorname{Ker}(\psi) = G/H$. By Claim 4.2.21 (ii), it follows that $\eta(\psi) = 1$. Thus Theorem A implies Claim 4.2.24. □

Claim 4.2.25. $e \neq 2$.

Proof. Assume that $e = 2$. Then $G/E \cong S_3$ or $G/E \cong \mathbf{Z}_3$ by Claim 4.2.19. By Claim 4.2.24 we have that there exists $L \lhd G$ such that $E < L$ and G/L is of the form (i), (ii), (iii), (iv) or (v). In this case, $G/L \cong \mathbf{Z}_2$ or $G = L$ since the only non-trivial normal subgroup of $G/E \cong S_3$ or $G/E \cong \mathbf{Z}_3$ is $\mathbf{Z}_3$. And $\mathbf{Z}_2$ is not among the possibilities given by Claim 4.2.24. So $e \neq 2$. □

Claim 4.2.26. $e \neq 3$.

Proof. Assume that $e = 3$. By Claim 4.2.19 we have that $G/E \cong \mathbf{Q}_8$ or $G/E \cong \mathrm{SL}(2,3)$. By Claim 4.2.24, we have that $E < L$, where L is a normal subgroup of G such that $\mathbf{F}(G/L)$ is isomorphic to either $\mathbf{Z}_3$ or $\mathbf{Q}_8$. This is impossible if $G/E \cong \mathbf{Q}_8$. Thus $G/E \cong \mathrm{SL}(2,3)$, $G/L \cong \mathbf{Z}_3$ and $\psi(1) = 2$. Let M be the normal subgroup of G such that $E < M$ and $M/E = \mathbf{F}(G/E) \cong Q_8$. By Claim 4.2.21 (ii) we have that $\alpha_1(1) = 3$ and $\alpha_1 \in \operatorname{Irr}(G \bmod E)$. Since $\alpha_1 \in \operatorname{Irr}(G \bmod E)$, $G/E \cong \mathrm{SL}(2,3)$ and $|G/M| = 3$, we have that α_1 is induced from a linear character of M. In particular

$$\alpha_1(g) = 0 \text{ for any } g \in G \setminus M. \tag{4.2.27}$$

Since $e = 3$, by Claim 4.2.20 we have that $\Lambda^e(1) = 3$. By Claim 4.2.21 (i) we have that $\alpha_2(1) = 8$. Since M is a normal subgroup of G, $(|G/M|, \alpha_2(1)) = 1$, and $\alpha_2 \in \operatorname{Irr}(G)$, by exercise 6.7 of [5] we have that $(\alpha_2)_M \in \operatorname{Irr}(M)$. Thus $(\alpha_2)_M \in \operatorname{Irr}(M \bmod Z)$, $(\alpha_2)_M(1) = 8$, E/Z is elementary abelian of order 9, $|M/E| = 8$ and E is a normal subgroup of G. By Clifford Theory we conclude that $(\alpha_2)_M$ is induced from a linear character of E. Therefore

$$\alpha_2(g) = 0 \text{ for any } g \in M \setminus E. \tag{4.2.28}$$

Since E is normal in G, while $(\alpha_2)_M \in \operatorname{Irr}(M)$ is induced from a linear character of E and $\alpha_2(1) = 8$, we have that $(\alpha_2)_E$ is the sum of 8 distinct linear characters of E. Since $\alpha_2 \in \operatorname{Irr}(G \bmod Z)$ and $|E/Z| = 9$ and $(\alpha_2)_E$ is the sum of 8 distinct linear characters of E, it follows that $(\alpha_2)_E = 1_Z^E - 1_E$. Therefore

$$\alpha_2(g) = -1 \text{ for any } g \in E \setminus Z. \tag{4.2.29}$$

By Claim 4.2.21 (iii) we must have that $\alpha_1\alpha_2 = 3\alpha_2$. By (4.2.27) and (4.2.28) we have that $\alpha_1\alpha_2(g) = 0$ for any $g \in G \setminus E$. Thus

$$\alpha_2(g) = 0 \text{ for any } g \in G \setminus E. \tag{4.2.30}$$

Therefore

$$\begin{aligned}
[\alpha_2,\alpha_2] &= \frac{1}{|G|}\sum_{g\in G}\alpha_2(g)\overline{\alpha_2(g)}\\
&= \frac{1}{|G|}\sum_{g\in E}\alpha_2(g)\overline{\alpha_2(g)} + \frac{1}{|G|}\sum_{g\in G\setminus E}\alpha_2(g)\overline{\alpha_2(g)}\\
&= \frac{1}{|G|}\sum_{g\in E}\alpha_2(g)\overline{\alpha_2(g)} \quad \text{by (4.2.30)}\\
&= \frac{1}{|G|}\sum_{g\in Z}\alpha_2(g)\overline{\alpha_2(g)} + \frac{1}{|G|}\sum_{g\in E\setminus Z}\alpha_2(g)\overline{\alpha_2(g)}\\
&= \frac{1}{|G|}|Z|\alpha_2(1)^2 + \frac{1}{|G|}(|E|-|Z|) \quad \text{because } \alpha_2\in \operatorname{Irr}(G \bmod Z) \text{ and (4.2.29)}\\
&= \frac{8^2}{|G/Z|} + \frac{(9-1)}{|G/Z|} \quad \text{since } \alpha_2(1)=8 \text{ and } |E/Z|=9\\
&= \frac{8\times 9}{24\times 9} = \frac{1}{3}.
\end{aligned}$$

Since $\alpha_2 \in \operatorname{Irr}(G)$, we have that $[\alpha_2,\alpha_2]=1$. But $[\alpha_2,\alpha_2]=\frac{1}{3}\neq 1$. Thus this case can not arise and $e\neq 3$. □

Claim 4.2.31. *If $e=5$ then $\psi(1)=2$, $\alpha_1(1)=3$, $\alpha_2(1)=24$, $m_2=4$ and $G/E\cong \mathrm{SL}(2,3)$. This gives case (i) in Proposition 4.2.13.*

Proof. By Claim 4.2.19 we have that $G/E\cong \mathrm{SL}(2,3)$. Analyzing the possibilities given by Claim 4.2.24, we have that $\psi(1)=2$. Thus by Claim 4.2.21 (ii) we have that $\alpha_1(1)=3$. Similarly, since $\Lambda^e(1)=5$, it follows by Claim 4.2.21 (i) that $\alpha_2(1)=24$ and by (4.2.22) that $\chi(1)=\psi(1)\Lambda^e(1)=10$. By Claim 4.2.21 (iii) we have that $m_2=1+\alpha_1(1)=1+3=4$ and we are done. □

Claim 4.2.32. *If $e=7$ then $\psi(1)=2$, $\alpha_1(1)=3$, $\alpha_2(1)=48$, $m_2=4$ and $G/E\cong \widetilde{\mathrm{GL}}(2,3)$. This gives case (ii) in Proposition 4.2.13.*

Proof. As in the proof of the previous claim, we have that $\psi(1)=2$ and $\alpha_1(1)=3$. Since $\Lambda^e(1)=7$, by Claim 4.2.21 (i) we have that $\alpha_2(1)=48$. Since $\Lambda^e(1)=\Lambda(1)=7$, by Claim 4.2.21 (ii) we have that $\alpha_2(1)=48$ and by (4.2.22) that $\chi(1)=\psi(1)\Lambda^e(1)=14$. By Claim 4.2.21 (iii) we have that $m_2=1+\alpha_1(1)=1+3=4$. □

It remains to analyze the case when $e=9$. We will use another approach to that case. The reason is that $H^2(E,\mathbf{C})\neq 1$ for an extra-special group E of order 2^5. Thus the character Λ may not extend to G. We start with some statements about χ, α_1 and α_2. Then we use that information to prove that $e=9$ can not hold.

Since $E\leq K$ and $K=\operatorname{Ker}(\alpha_1)$, the restriction χ_E is reducible. Since E/Z is a fully ramified section of G with respect to θ_E, we have that

$$\chi(1)=em \quad \text{where } m\geq 2. \tag{4.2.33}$$

Since E/Z is a fully ramified section of G with respect to θ_E and $Z=\mathbf{Z}(G)$, $\chi(g)=0$ for all $g\in E\setminus Z$. Thus $\chi\overline{\chi}(g)=0=1+\alpha_1(1)+m_2\alpha_2(g)$ for all $g\in E\setminus Z$.

Therefore for all $g \in E \setminus Z$ we have

$$\alpha_2(g) = -\frac{1+\alpha_1(1)}{m_2}. \tag{4.2.34}$$

Recall that $\alpha_2 \in \mathrm{Irr}(G)$. So $\alpha_2(g)$ is an algebraic integer for any $g \in G$. Since $\frac{1+\alpha_1(1)}{m_2}$ is a rational algebraic integer, it is an integer. Thus we conclude that

$$m_2 \text{ divides } 1+\alpha_1(1). \tag{4.2.35}$$

By (4.1.2) we have that

$$\alpha_2(1) = \frac{\chi(1)^2}{m_2} - \frac{1+\alpha_1(1)}{m_2}. \tag{4.2.36}$$

By (4.2.34) and (4.2.36) we have that

$$(\alpha_2)_E = s\,1_Z^E - \frac{1+\alpha_1(1)}{m_2} 1_E$$

for some integer $s \geq \frac{1+\alpha_1(1)}{m_2}$. So

$$se^2 - \frac{1+\alpha_1(1)}{m_2} = \alpha_2(1) = \frac{m^2e^2}{m_2} - \frac{1+\alpha_1(1)}{m_2}.$$

Hence $s = m^2/m_2$. Therefore

$$m_2 \text{ divides } m^2. \tag{4.2.37}$$

Since $\alpha_2 \in \mathrm{Irr}(G)$ and $\mathrm{Ker}(\alpha_2) = Z$, we have that

$$\alpha_2(1)^2 < |G:Z| \text{ and } \alpha_2(1) \text{ divides } |G:Z|. \tag{4.2.38}$$

Claim 4.2.39. *Assume $e = 9$. Then $E < K = \mathrm{Ker}(\alpha_1)$. In particular, α_1 is not a faithful character of $G \bmod E$.*

Proof. Suppose $E = K$. Observe that $\mathbf{Z}(\mathbf{F}(\bar{G}))$ is a non-trivial cyclic group by Claim 4.2.19 (v). Therefore there exists a cyclic chief factor H/E of G with $H/E \leq \mathbf{Z}(\mathbf{F}(\bar{G}))$. Since $\Lambda \in \mathrm{Irr}(E)$ is G-invariant and H/E is cyclic, Λ extends to H. We are assuming that $E = K = \mathrm{Ker}(\alpha_1)$. Thus $H \not\leq K$, and $H \not\leq \mathrm{Ker}(\alpha_2)$. Hence $\chi_H \in \mathrm{Irr}(H)$ by Lemma 4.2.1. Since Λ extends to H, H/Z is cyclic and $[\chi_E, \Lambda] \neq 0$, it follows that $\chi_E = \Lambda$. By Claim 4.2.20 we have that $\Lambda(1) = e$. Thus $\chi(1) = e$. But that is a contradiction with (4.2.33). □

Claim 4.2.40. $e \neq 9$.

Proof. Assume that $e = 9$. By Claim 4.2.19 (v) we have that

$$|G:Z| \text{ divides } 10 \times 32 \times 81 = 25920. \tag{4.2.41}$$

By Claim 4.2.17, Claim 4.2.19 and Lemma 4.2.12 we have that $\alpha_1(1) \in \{1,2,4,5,8,10\}$. By Claim 4.2.39 and Lemma 4.2.12 we have that $\alpha_1(1) \neq 8$. Thus

$$\alpha_1(1) \in \{1,2,4,5,10\}. \tag{4.2.42}$$

This and (4.2.35) imply that

$$m_2 \in \{1,2,3,5,6,11\}. \tag{4.2.43}$$

Therefore

$$\frac{1+\alpha_1(1)}{m_2} \leq 11. \tag{4.2.44}$$

If $m_2 = 1$, then by (4.2.33) and (4.2.36) we have that

$$\begin{aligned}\alpha_2(1) &= \frac{\chi(1)^2}{m_2} - \frac{1+\alpha_1(1)}{m_2} \\ &\geq 4 \times 81 - 11 = 313.\end{aligned}$$

But then $\alpha_2(1)^2 \geq 313^2 = 97969 > 25920 \geq |G : Z|$. This is impossible by (4.2.38). So $m_2 \neq 1$.

If $m_2 = 2$, then by (4.2.35) we have that $\alpha_1(1) = 1$ or $\alpha_1(1) = 5$. By (4.2.37) we have that $2 \mid m$ If $m > 2$ then $m \geq 4$ and

$$\frac{\chi(1)^2}{m_2} = \frac{m^2 9^2}{2} \geq \frac{4^2 \times 9^2}{2} = 8 \times 81.$$

Thus by (4.2.36) we have that $\alpha_2(1) \geq 8 \times 81 - 3$ and therefore $\alpha_2(1)^2 > |G : Z|$. So we may assume that $m = 2$. Since $\alpha_1(1) \in \{1, 5\}$ and $m_2 = 2$, by (4.2.36) either $\alpha_2(1) = 2 \times 81 - 1 = 161$ or $\alpha_2(1) = 2 \times 81 - 3 = 159$. But neither 161 nor 159 divides $|G : Z|$. Hence $m_2 \neq 2$.

Suppose $m_2 \in \{3, 5, 6, 11\}$. By (4.2.37) we have that $m_2 \mid m^2$. Thus $m \geq 9$. Therefore

$$\frac{\chi(1)^2}{m_2} \geq \frac{9^2 9^2}{3} = 2187.$$

Thus $\alpha_2(1) \geq 2187 - 11$ by (4.2.44) and (4.2.36). By (4.2.41) $\alpha_2(1)^2 = 2176^2 > 25920 \geq |G : Z|$, a contradiction with (4.2.38). We conclude that $m_2 \notin \{3, 5, 6, 11\}$.

We have considered all the possible options given by (4.2.43) and we have showed that none of them is possible. We conclude that $e \neq 9$. □

We have analyzed all the options given by Claim 4.2.19, so the proof is complete. □

We summarize some of the information we obtained in the previous lemmas.

Theorem B. *Assume Hypothesis 4.1.1. Then*

(i) The order of G is even.

(ii) $\mathrm{dl}(G) \leq 18$.

(iii) Either $\mathrm{Ker}(\alpha_1) = Z$ *or* $\mathrm{Ker}(\alpha_2) = Z$. *If* $\mathrm{Ker}(\alpha_1)$ *and* $\mathrm{Ker}(\alpha_2)$ *are both abelian subgroups, then $\chi(1)$ is a power of a prime number. Otherwise* $\{m_1, m_2\} = \{1, 4\}$ *and* $\chi(1) \in \{10, 14\}$.

Proof. The order of G is even by Proposition 4.2.8. Either $\mathrm{Ker}(\alpha_1) = Z$ or $\mathrm{Ker}(\alpha_2) = Z$ by Lemma 4.2.4. Propositions 4.2.5, 4.2.9 and 4.2.13 imply (ii) and (iii). □

4.3. **Examples.** In this subsection, we study examples of groups satisfying Hypothesis 4.1.1.

Lemma 4.3.1. *Let G be a finite group and E be a nonabelian subgroup. Assume $E/\mathbf{Z}(G)$ is an abelian chief factor of G and G/E acts on $(E/\mathbf{Z}(G))^{\#}$ with 2 orbits. Let $\chi \in \mathrm{Irr}(G)$ be a faithful character. Assume also that $\chi_E \in \mathrm{Irr}(E)$. Then*

$$\chi\overline{\chi} = 1_G + \alpha_1 + \alpha_2$$

where α_1 and α_2 are distinct irreducible characters of G and $\mathrm{Ker}(\alpha_1) = \mathrm{Ker}(\alpha_2) = \mathbf{Z}(G)$.

Proof. Set $Z = \mathbf{Z}(G)$. Let $\lambda \in \operatorname{Irr}(Z)$ be such that $[\lambda, \chi_Z] \neq 0$. Since $\chi_E \in \operatorname{Irr}(E)$ is a faithful character of E and E is not abelian, we have that $\chi(1) > 1$. Thus χ_Z is a reducible character. Since $\chi_E \in \operatorname{Irr}(E)$, E/Z is an abelian chief factor, χ_Z is a reducible character and $Z = \mathbf{Z}(G)$, we have that E/Z is a fully ramified section with respect to χ_E and λ, and that $\chi(1)^2 = |E : Z|$. Thus

$$(\chi\overline{\chi})_E = 1_Z^G. \tag{4.3.2}$$

Since G acts on $(E/\mathbf{Z}(G))^\#$ with 2 orbits, G acts on $\operatorname{Irr}(E \bmod Z)^\#$ with 2 orbits.

By Clifford Theory we have that for any character $\alpha \in \operatorname{Irr}(G)$, the irreducible constituents of α_E form a G-orbit. Let $\alpha_1 \in \operatorname{Irr}(G)^\#$ be a character such that $[\alpha_1, \chi\overline{\chi}] = m_1 \neq 0$. By Lemma 3.1.2 we have that $\operatorname{Ker}(\alpha_1) \geq Z$. Thus the irreducible constituents of $(\alpha_1)_E$ lie in $\operatorname{Irr}(E \bmod Z)^\#$. Since G acts on $\operatorname{Irr}(E \bmod Z)^\#$ with 2 orbits and the irreducible constituents of $(\alpha_1)_E$ form a G-orbit, there exists $\delta \in \operatorname{Irr}(E \bmod Z)$ such that $[\delta, (\alpha_1)_E] = 0$. Then there exists a character $\alpha_2 \in \operatorname{Irr}(G)$ such that $[(\alpha_2)_E, \delta] \neq 0$ and $[\alpha_2, \chi\overline{\chi}] = m_2 \neq 0$. Thus the irreducible constituents of $(\alpha_1)_E$ and $(\alpha_2)_E$ form two distinct G-orbits of $\operatorname{Irr}(E \bmod Z)^\#$. Since G acts on $\operatorname{Irr}(E \bmod Z)^\#$ with 2 orbits we conclude that those are the only distinct G-orbits of $\operatorname{Irr}(E \bmod Z)^\#$.

Since E/Z is an abelian chief factor of G, by (4.3.2) we have that the irreducible constituents of $\chi\overline{\chi}$ appear with multiplicity 1. Since the irreducible constituents of $(\alpha_1)_E$ and $(\alpha_2)_E$ form two distinct G-orbits of $\operatorname{Irr}(E \bmod Z)^\#$ and the irreducible constituents of $\chi\overline{\chi}$ appear with multiplicity 1, we have that α_1 and α_2 are all the non-principal irreducible constituents of $\chi\overline{\chi}$ and they appear with multiplicity 1. Thus

$$\chi\overline{\chi} = 1 + \alpha_1 + \alpha_2.$$

□

Proposition 4.3.3. *There exist a group G and a character $\chi \in \operatorname{Irr}(G)$ such that Hypothesis 4.1.1 holds and*

$$\operatorname{Ker}(\alpha_1) = \operatorname{Ker}(\alpha_2).$$

Thus there exist a group G and a character $\chi \in \operatorname{Irr}(G)$ satisfying the hypotheses of Proposition 4.2.5.

Proof. Let E be an extraspecial group of order 27 and exponent 3. The 2-Sylow subgroup of $\operatorname{SL}(2,3)$ is isomorphic to the group $\mathbf{Q}_8$ by Lemma 2.1.3. Thus $\operatorname{SL}(2,3)$ contains a cyclic group of order 4. Let H be a cyclic group of order 4. By Lemma 2.1.3 (i) and Lemma 3.3.1, we may assume that H is a subgroup of $\operatorname{Aut}(E \bmod \mathbf{Z}(E))$. We can check that H acts on $(E/Z)^\#$ with 2 orbits of the same size.

Set $G = E \rtimes H$. Observe that $E/\mathbf{Z}(E)$ is an abelian chief factor of G. Also observe that G/E acts faithfully on $E/\mathbf{Z}(E)$, and with 2 orbits on $(E/\mathbf{Z}(E))^\#$. Let $\Lambda \in \operatorname{Irr}(E)$ be a character of degree 3. Observe that Λ is G-invariant. Since $(|H|, |E|) = 1$ and Λ is G-invariant, we have that Λ extends to G. Let $\chi \in \operatorname{Irr}(E)$ be an extension of Λ. Observe that the hypotheses of Lemma 4.3.1 hold for G and χ. Thus the result follows from Lemma 4.3.1. □

Proposition 4.3.4. *Let p be a prime number and $m > 0$ an integer. Then there exist a solvable group G and a character $\chi \in \operatorname{Irr}(G)$ such that $\chi(1) = p^m$ and the hypotheses of Proposition 4.2.9 hold. Also, there exist a supersolvable group G and a character $\chi \in \operatorname{Irr}(G)$ such that $\chi(1) = p$ and the hypotheses of Proposition 4.2.9 hold.*

Proof. Let $\mathrm{GF}(q)$ be a finite field with q elements, where $q = p^m$. Denote by F^* the group of units of $\mathrm{GF}(q)$. Also denote by F the additive group of $\mathrm{GF}(q)$. Observe that F^* acts on F by multiplication.

Define M to be the semi-direct product $F \rtimes F^*$ of F by F^*. Thus, for any $f_1, g_1 \in F$ and $f_2, g_2 \in F^*$, we have

$$(g_1 \rtimes g_2)(f_1 \rtimes f_2) = (g_1 + g_2 f_1) \rtimes g_2 f_2.$$

Observe that if $m = 1$ the group M is a cyclic by cyclic group and thus is supersolvable.

Set $X = \mathrm{GF}(q)$. For any $f_1 \rtimes f_2 \in M$ and $x \in X$, define

$$(f_1 \rtimes f_2)(x) = f_1 + f_2 x.$$

We can check that this defines an action of M on the set X. We can also check that $0 \in X$ is fixed by F^*.

Let C be a cyclic group of order p. Set

$$K = C^X = \text{ the set of all functions from } X \text{ to } C.$$

Observe that K is just an elementary abelian group of order $p^{|X|}$. We can check that

$$(fm)(x) = f(mx),$$

for any $m \in M$ and $x \in X$, defines an action of M on K.

Let G be the wreath product of C and M relative to X, i.e. $G = K \rtimes M$. Observe that if $m = 1$, then the group G is a supersolvable group.

Let $\lambda \in \mathrm{Irr}(C)^{\#}$ be a character. Choose $\theta_0 \in \mathrm{Irr}(K)$ so that we have, for all $k \in K$,

$$\theta_0(k) = \lambda(k(0)), \text{ where } 0 = 0_F \in X.$$

Since $0 = 0_F \in X$ is fixed by F^*, the stabilizer of θ_0 in G is $G_{\theta_0} = K \rtimes F^*$. We can check that θ_0 extends to G_{θ_0}. Let ${\theta_0}^e \in \mathrm{Irr}(G_{\theta_0})$ be an extension of θ_0. Set $(\theta_0^e)^G = \chi$. By Clifford theory we have that $\chi \in \mathrm{Irr}(G)$. Observe that $\chi(1) = q$ since $G_{\theta_0} = F^*K$ and θ_0 extends to G_{θ_0}. Since ${\theta_0}^e \in \mathrm{Irr}(G_{\theta_0})$ is an extension of θ_0 and $(\theta_0^e)^G = \chi$, we have that $[\chi_K, \theta_0] = 1$. We can check that

$$\chi_K = \sum_{x \in X} \theta_x$$

where $\theta_x(k) = \lambda(k(x))$ for all $k \in K$. Therefore

$$(\chi\overline{\chi})_K = \sum_{x,y \in X} \theta_x \overline{\theta_y}.$$

Since $G_{\theta_0} = K \rtimes F^*$, $G = (K \rtimes F^*)(K \rtimes F)$ and $(K \rtimes F^*) \cap (K \rtimes F) = K$, we have that $\chi_{K \rtimes F} = (\theta_0)^{K \rtimes F} \in \mathrm{Irr}(K \rtimes F)$. Thus $\chi\overline{\chi}(g) = 0$ for all $g \in (K \rtimes F) \setminus K$. We can check that for any $\gamma \in \mathrm{Irr}((K \rtimes F) \bmod K)$,

$$[\gamma, (\chi\overline{\chi})_{K \rtimes F}] \neq 0. \tag{4.3.5}$$

We can check that $\alpha_1 = \gamma^G \in \mathrm{Irr}(G)$. Since $|G : K \rtimes F| = |F^*| = q - 1$, we have that $\alpha_1(1) = q - 1$. Observe that (4.3.5) implies

$$[\alpha_1, \chi\overline{\chi}] \neq 0. \tag{4.3.6}$$

Define $\delta = \theta_0\overline{\theta_1}$. Observe that $\theta_x(1) = 1$ for all $x \in X$. Therefore $\delta \in \mathrm{Irr}(K)$. Since 0 and 1 are not fixed simultaneously by any non-trivial element of M, we have that $G_\delta = K$. Thus $\delta^G = \alpha_2 \in \mathrm{Irr}(G)$. Observe that

$$[\alpha_2, \chi\overline{\chi}] \neq 0 \tag{4.3.7}$$

since $[\delta, (\chi\overline{\chi})_K] \neq 0$. Observe that $\alpha_2(1) = q(q-1) = |G : K|$.

Observe that $\chi(1) = q$, $\alpha_1(1) = q-1$, $\alpha_2(1) = q(q-1)$, (4.3.6) and (4.3.7) imply

$$\chi\overline{\chi} = 1_G + \alpha_1 + \alpha_2.$$

□

Proposition 4.3.8. *There exist a solvable group G and a faithful character $\chi \in \mathrm{Irr}(G)$ such that $\chi(1) = 10$ and*

$$\chi\overline{\chi} = 1_G + \alpha_1 + 4\alpha_2,$$

where $\alpha_1, \alpha_2 \in \mathrm{Irr}(G)$, $\alpha_1(1) = 3$ and $\alpha_2(1) = 24$. Thus there exist a group G and a character $\chi \in \mathrm{Irr}(G)$ satisfying the conditions in case (i) of Proposition 4.2.13.

Proof. Let E be an extra-special group of order 5^3 and exponent 5. Then $E/\mathbf{Z}(E)$ is an elementary abelian group of order 25. By Lemma 2.1.3 (iv) there exists a subgroup H of $\mathrm{SL}(2,5)$ isomorphic to $\mathrm{SL}(2,3)$. Assume H acts on the vector space $E/\mathbf{Z}(E)$ in the natural way. We can check that in fact H acts on $(E/\mathbf{Z}(E))^\#$ regularly and transitively. By Lemma 2.1.3 (i), we have that $\mathrm{Sp}(2,5) \cong \mathrm{SL}(2,5)$. By Lemma 3.3.1, there exists a exact sequence

$$0 \to \mathrm{Inn}(E) \to \mathrm{Aut}(E \bmod \mathbf{Z}(E)) \to \mathrm{Sp}(2,5) \to 0.$$

Since $\mathrm{Sp}(2,5) \cong \mathrm{SL}(2,5)$, we may assume that H acts on E. Define $G = E \rtimes H$ to be the semi-direct product of E by H. Then $\mathbf{Z}(G) = \mathbf{Z}(E)$.

Let $\Lambda \in \mathrm{Irr}(E)$ be a non-linear character. Observe that Λ is G-invariant. Since H acts on E with coprime action, Λ extends to G, i.e, there exists $\phi \in \mathrm{Irr}(G)$ such that $\phi_E = \Lambda$. Let $\gamma \in \mathrm{Irr}(G \bmod E)$ be a character of degree 2. Set $\chi = \phi\gamma$. Then $\chi \in \mathrm{Irr}(G)$ and $\chi(1) = 10$.

Observe that $\chi(e) = 0$ for any $e \in E \setminus \mathbf{Z}(E)$.

Let $\delta \in (\mathrm{Lin}(E \bmod \mathbf{Z}))^\#$. Since H acts on $(E/\mathbf{Z}(E))^\#$ regularly and transitively, δ induces irreducibly. Let $\alpha_2 = \delta^G \in \mathrm{Irr}(G)$. Observe that $\alpha_2(1) = 24$. Also $\alpha_2(g) = 0$ if $g \in G \setminus E$. Thus

$$\begin{aligned}
[\chi\overline{\chi}, \alpha_2] &= \frac{1}{|G|}\sum_{g\in G}(\chi\overline{\chi})(g)\ \overline{\alpha_2(g)} \\
&= \frac{1}{|G|}[\sum_{g\in G\setminus E}(\chi\overline{\chi})(g)\ \overline{\alpha_2(g)} + \sum_{g\in E\setminus \mathbf{Z}(G)}(\chi\overline{\chi})(g)\ \overline{\alpha_2(g)} + \sum_{g\in \mathbf{Z}(G)}(\chi\overline{\chi})(g)\ \overline{\alpha_2(g)}] \\
&= \frac{1}{|G|}\sum_{g\in \mathbf{Z}(G)}(\chi\overline{\chi})(g)\ \overline{\alpha_2(g)} = 4.
\end{aligned}$$

We can check that $\gamma\overline{\gamma} = 1 + \alpha_1$, where $\alpha_1(1) = 3$ and $\alpha_1 \in \mathrm{Irr}(G \bmod E)$. Thus

$$\begin{aligned}
[\chi\overline{\chi}, \alpha_1] &= [\chi, \chi\alpha_1] \\
&= [\phi\gamma, \phi\gamma\alpha_1] \\
&= [\phi\gamma\overline{\gamma}, \phi\alpha_1] = 1
\end{aligned}$$

where the last inequality is since $[\gamma\overline{\gamma}, \alpha_1] = 1$.

Thus $\chi\overline{\chi} = 1_G + \alpha_1 + 4\alpha_2$, since $(\chi\overline{\chi})(1) = 10^2 = 1+3+96 = 1+\alpha_1(1)+\alpha_2(1)$. $\square$

Proposition 4.3.9. *There exist a solvable group G and a faithful character $\chi \in \mathrm{Irr}(G)$ such that*

$$\chi\overline{\chi} = 1_G + \alpha_1 + 4\alpha_2$$

where $\alpha_1, \alpha_2 \in \mathrm{Irr}(G)$, $\chi(1) = 14$, $\alpha_1(1) = 3$ and $\alpha_2(1) = 48$. Thus there exist a group G and a character $\chi \in \mathrm{Irr}(G)$ satisfying the conditions in case (ii) of Proposition 4.2.13.

Proof. Let E be an extra-special group of order 7^3 and exponent 7. Observe that $E/\mathbf{Z}(E)$ is an elementary abelian 7-group. By Lemma 2.1.6, we may regard $\widetilde{\mathrm{GL}}(2,3)$ as a subgroup of $\mathrm{SL}(2,7)$. By Lemma 2.1.3 and Lemma 3.3.1, we may assume that $\widetilde{\mathrm{GL}}(2,3)$ acts on E. Define G to be the semi-direct product $E \rtimes \widetilde{\mathrm{GL}}(2,3)$ of E by $\widetilde{\mathrm{GL}}(2,3)$. We can check as before that there exists a faithful character $\chi \in \mathrm{Irr}(G)$ satisfying the conditions of the proposition. $\square$

4.4. **Dihedral and nilpotent groups.** Using the character table of the dihedral group of order 2^m with $m > 3$, we can check that dihedral groups have characters χ that satisfy Hypothesis 4.1.1. Observe that therefore the index $|G : \mathbf{Z}(G)|$ is not going to be bounded as it was in the case where $\chi\overline{\chi} = 1_G + m\alpha$. We are going to see now that this is, in some sense, the general case whenever we are working with nilpotent groups.

Theorem 4.4.1. *Assume Hypothesis 4.1.1. Assume also that G is nilpotent. Then*

$$m_1 = m_2 = 1, \quad \chi(1) = 2$$

and $G/\mathbf{Z}(G)$ is isomorphic to a dihedral 2-group.

Proof. By Lemma 4.2.4, either $\mathrm{Ker}(\alpha_1) = Z$ or $\mathrm{Ker}(\alpha_2) = Z$. Since the chief factors of a nilpotent group are cyclic of prime order, we have by Propositions 4.2.5, 4.2.13 and 4.2.9 that either $\mathrm{Ker}(\alpha_1) \neq Z$ and $\mathrm{Ker}(\alpha_1)$ is abelian, or $\mathrm{Ker}(\alpha_2) \neq Z$ and $\mathrm{Ker}(\alpha_2)$ is abelian. Without lost of generality we may assume that $\mathrm{Ker}(\alpha_1) \neq Z$ is an abelian group and $\mathrm{Ker}(\alpha_2) = Z$.

We use now the notation of Lemma 4.2.9. Then $|G : G_\theta| = |L : K|$ is a prime number. Since G is nilpotent, G_θ is a normal subgroup of G. Observe that $K \leq G_\theta$. If $G_\theta \neq K$, then $\chi_{G_\theta} \in \mathrm{Irr}(G_\theta)$ by Lemma 4.2.1. But this can not be since $\chi = \chi_\theta^G$ and $G_\theta < G$. Thus $G_\theta = K$ and so $G = L$.

Since $K = \mathrm{Ker}(\alpha_1)$ is abelian and $|G : K| = p$ is a prime number, $\alpha_1(1) = 1$ and either $\alpha_2(1) = 1$ or $\alpha_2(1) = p$. Assume that $\alpha_2(1) = 1$. Since $\mathrm{Ker}(\alpha_2) = Z$, the group G/Z is cyclic. Since $Z = \mathbf{Z}(G)$ and G/Z is cyclic, it follows that G is an abelian group. Thus $\chi(1) = 1$. But then χ can not satisfy (4.1.2). So $\alpha_2(1) \neq 1$. We conclude that $\alpha_2(1) = p$.

Since χ and α_2 are induced from characters of K, we have $\chi(g) = \alpha_2(g) = 0$ for all $g \in G \setminus K$. Thus $\alpha_1(g) = -1/a_1 = -1$ for all $g \in G \setminus K$. Since G/K is a cyclic group of order p, $\alpha_1(g) = -1$ for all $g \in G \setminus K$ and $\alpha_1(1) = 1$, we must have that $p = 2$. Since $\chi(1) = p$, $\alpha_2(1) = p$ and $\chi(1)^2 = 1 + \alpha_1(1) + m_2\alpha_2(1)$, we conclude that $m_2 = 1$.

We want to see now that G/Z is isomorphic to a dihedral 2-group. Let $\chi_K = \theta_1 + \theta_2$ and $(\alpha_2)_K = \gamma_1 + \gamma_2$, where $\theta_1, \theta_2, \gamma_1, \gamma_2 \in \mathrm{Irr}(K)$, $\theta_1 \neq \theta_2$ and $\gamma_1 \neq \gamma_2$.

This holds because χ and α_2 are induced from characters of the abelian normal subgroup K and $|G : K| = 2$. Then by (4.1.2) we have that

$$\begin{aligned}\chi_K\overline{\chi_K} &= (\theta_1+\theta_2)\overline{(\theta_1+\theta_2)}\\ &= 2\,1_K + \theta_1\overline{\theta_2} + \overline{\theta_1}\theta_2\\ &= 2\,1_K + (\alpha_2)_K\\ &= 2\,1_K + \gamma_1 + \gamma_2.\end{aligned}$$

Choose $\gamma_1 = \theta_1\overline{\theta_2}$ and $\gamma_2 = \overline{\theta_1}\theta_2$. Observe that $\theta_1\overline{\theta_2} = \overline{\overline{\theta_1}\theta_2}$. Thus $\gamma_1 = \overline{\gamma_2}$. If γ_1 is a real character, then $\gamma_1 = \overline{\gamma_1} = \gamma_2$. But $\gamma_1 \neq \gamma_2$. Therefore γ_1 and γ_2 are not real characters.

Since γ_1 is a linear character, it follows that $(\gamma_1)^{-1} = \overline{\gamma_1} = \gamma_2$. In particular $\mathrm{Ker}(\gamma_1) = \mathrm{Ker}(\gamma_2)$. Therefore $Z = \mathrm{Ker}(\alpha_2) = \mathrm{Ker}(\gamma_1) \cap \mathrm{Ker}(\gamma_2) = \mathrm{Ker}(\gamma_1)$. Since $\gamma_1(1) = 1$, the group K/Z is cyclic. Let $\bar{x}$ be a generator of K/Z. Let y be a coset representative of $G \setminus K$. Since $(\theta)_K = \gamma_1 + \gamma_2$, $\gamma_1 \neq \gamma_2$ and y is a coset representative of $G\backslash K$, by Clifford theory it follows that $(\gamma_1)^y = \gamma_2$. Since $\gamma_1^{y^2} = \gamma_1$, $\gamma_1^y = \gamma_2 = (\gamma_1)^{-1}$ and $\gamma_1 \in \mathrm{Irr}(K \bmod Z)$ is a faithful character, we have that

$$\bar{y}\bar{x}\bar{y}^{-1} = \bar{x}^{-1}. \tag{4.4.2}$$

Thus G/Z is either isomorphic to the dihedral group or to the generalized quaternion group. If G/Z is isomorphic to the generalized quaternion group, then G is a non-split central extension of a cyclic group by a generalized quaternion group. But that can not be since $H^2(\mathbf{Q}_{2^n}, \mathbf{C}) = 1$ for any $n \geq 3$ (see example in page 301 of [2]). Thus G/Z is isomorphic to the dihedral 2-group and the proof is complete. □

REFERENCES

[1] E. Adan-Bante, Products of characters and derived length, J. Algebra 266 (2003), 305-319.

[2] C. W. Curtis, I. Reiner, Methods of Representation Theory Vol 1 with applications to finite groups and orders, John Wiley & Sons Inc, New York 1990.

[3] D. Gorenstein, Finite Groups. New York: Harper and Row 1968.

[4] B. Huppert, Endliche Gruppen I, Berlin–Heidelberg–New York : Springer 1967.

[5] I. M. Isaacs, Character Theory of Finite Groups. New York-San Francisco–London: Academic Press 1976.

[6] T. M. Keller, Orbit sizes and character degrees II, J. reine angew. Math. 516 (1999), 27-114.

[7] D. J. S. Robinson, A Course in the Theory of Groups, Springer-Verlag, New York 1996.

[8] D. L. Winter, The automorphism group of an extraspecial p-group, Rocky mountain J. of Math 2 (1972), 159-168.

[9] O. Manz, T. R. Wolf, Representations of Solvable Groups, Lecture Notes Series Vol 185, Cambridge University Press 1993.

UNIVERSITY OF SOUTHERN MISSISSIPPI AT GULF COAST, LONG BEACH, MS, 39560
E-mail address: `Edith.Bante@usm.edu`